

Patent Application

**MAMMALIAN RECEPTOR PROTEINS;  
RELATED REAGENTS AND METHODS**

INVENTORS:

Madaline Chirica, a citizen of the United States, residing at 1520 University Avenue,  
Palo Alto, California 94301;

Robert A. Kastelein, a citizen the Netherlands, residing at 463 Summit Drive, Redwood  
City, California 94062;

Kevin W. Moore, a citizen of the United States residing at 251 Carolina Lane; Palo Alto,  
California 94306; and


Christi L. Parham; a citizen of the United States, residing at 2385 30th Avenue; San  
Francisco, California 94116.

"Express Mailing" mailing label number EL 982 851 390 US

Date of Deposit September 18, 2003

I hereby certify that this paper is being deposited with the United States Postal Service

"Express Mail Post to Addressee" service under 37 C.F.R. §1.10 on the date indicated above  
and is addressed to: Mail Stop Patent Application, Commissioner for Patents; P.O. Box 1450,  
Alexandria, VA 22313-1450.

  
Melanie Lyons

9-18-03  
Date

Assignee:

Schering Corporation, a New Jersey corporation.

**Customer No. 28008**

DNAX Research Institute  
901 California Avenue  
Palo Alto, CA 94304-1104  
Tel: (650) 496-1244  
Fax: (650) 496-1200

## MAMMALIAN RECEPTOR PROTEINS; RELATED REAGENTS AND METHODS

This filing claims benefit of U.S. Provisional Patent Application no. 60/203,426, filed May 10, 2000 which is incorporated herein by reference in its entirety.

5

### FIELD OF THE INVENTION

The present invention relates to compositions and methods for affecting mammalian physiology, including immune system function. In particular, it provides methods to regulate development and/or the immune system. Diagnostic and therapeutic uses of these materials are also disclosed.

10

### BACKGROUND OF THE INVENTION

Recombinant DNA technology refers generally to techniques of integrating genetic information from a donor source into vectors for subsequent processing, such as through introduction into a host, whereby the transferred genetic information is copied and/or expressed in the new environment. Commonly, the genetic information exists in the form of complementary DNA (cDNA) derived from messenger RNA (mRNA) coding for a desired protein product. The carrier is frequently a plasmid having the capacity to incorporate cDNA for later replication in a host and, in some cases, actually to control expression of the cDNA and thereby direct synthesis of the encoded product in the host. See, e.g., Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.) vols. 1-3, CSH Press, NY.

For some time, it has been known that the mammalian immune response is based on a series of complex cellular interactions, called the "immune network". Recent research has provided new insights into the inner workings of this network. While it remains clear that much of the immune response does, in fact, revolve around the network-like interactions of lymphocytes, macrophages, granulocytes, and other cells, immunologists now generally hold the opinion that soluble proteins, known as lymphokines, cytokines, or monokines, play critical roles in controlling these cellular interactions. Thus, there is considerable interest in the isolation, characterization, and mechanisms of action of cell modulatory factors, an understanding of which will lead to significant advancements in the diagnosis and therapy of numerous medical abnormalities, e.g., immune system disorders.

Lymphokines apparently mediate cellular activities in a variety of ways. See, e.g., Paul (ed. 1996) Fundamental Immunology 3d ed., Raven Press, New York; and Thomson (ed. 1994) The Cytokine Handbook 2d ed., Academic Press, San Diego. They have been shown to support the proliferation, growth, and/or differentiation of pluripotential hematopoietic stem cells into vast numbers of progenitors comprising diverse cellular lineages which make up a complex immune system. Proper and balanced interactions between the cellular components

35

are necessary for a healthy immune response. The different cellular lineages often respond in a different manner when lymphokines are administered in conjunction with other agents.

Cell lineages especially important to the immune response include two classes of lymphocytes: B-cells, which can produce and secrete immunoglobulins (proteins with the capability of recognizing and binding to foreign matter to effect its removal), and T-cells of various subsets that secrete lymphokines and induce or suppress the B-cells and various other cells (including other T-cells) making up the immune network. These lymphocytes interact with many other cell types.

Research to better understand and treat various immune disorders has been hampered by the general inability to maintain cells of the immune system in vitro. Immunologists have discovered that culturing many of these cells can be accomplished through the use of T-cell and other cell supernatants, which contain various growth factors, including many of the lymphokines.

Various growth and regulatory factors exist which modulate morphogenetic development. Many receptors for cytokines are known. Often, there are at least two critical subunits in the functional receptor. See, e.g., Heinrich, et al. (1998) *Biochem. J.* 334:297-314; Gonda and D'Andrea (1997) *Blood* 89:355-369; Presky, et al. (1996) *Proc. Nat'l Acad. Sci. USA* 93:14002-14007; Drachman and Kaushansky (1995) *Curr. Opin. Hematol.* 2:22-28; Theze (1994) *Eur. Cytokine Netw.* 5:353-368; and Lemmon and Schlessinger (1994) *Trends Biochem. Sci.* 19:459-463.

From the foregoing, it is evident that the discovery and development of new soluble proteins and their receptors, including ones similar to lymphokines, should contribute to new therapies for a wide range of degenerative or abnormal conditions which directly or indirectly involve development, differentiation, or function, e.g., of the immune system and/or hematopoietic cells. In particular, the discovery and understanding of novel receptors for lymphokine-like molecules which enhance or potentiate the beneficial activities of other lymphokines would be highly advantageous. The present invention provides new receptors for ligands exhibiting similarity to cytokine like compositions and related compounds, and methods for their use.

30

## SUMMARY OF THE INVENTION

The present invention is directed to novel receptors related to cytokine receptors, e.g., primate, cytokine receptor-like molecular structures, designated DNAX Cytokine Receptor Subunits (DCRS), and their biological activities. In particular, it provides description of one subunit, designated DCRS5. It includes nucleic acids coding for the polypeptides themselves and methods for their production and use. The nucleic acids of the invention are characterized,

35

in part, by their homology to cloned complementary DNA (cDNA) sequences enclosed herein. Additionally, the invention provides matching of the p40/IL-B30 ligand with receptor subunits DCRS5 and IL-12R $\beta$ 1, which pairing provides insight into indications for use of the agonists and antagonists based upon reagents directed thereto.

5       The present invention provides a substantially pure or recombinant polypeptide comprising at least ten contiguous amino acids of the intracellular portion of SEQ ID NO: 2. In certain embodiments, the polypeptide: comprises at least 25 contiguous amino acids of the intracellular portion of SEQ ID NO: 2; is recombinant, comprising the intracellular portion of SEQ ID NO: 2; further comprises at least ten contiguous amino acids of the non-intracellular  
10   portion of SEQ ID NO: 2; comprises at least 25 amino acids of the extracellular portion of SEQ ID NO: 2; comprises the mature SEQ ID NO: 2; or is a substantially pure natural polypeptide. In others, the recombinant polypeptide: consists of the mature sequence of Table 1; is an unglycosylated polypeptide; is from a human; comprises at least 40 contiguous amino acids of SEQ ID NO: 2; exhibits at least three nonoverlapping segments of at least fifteen contiguous  
15   amino acids of SEQ ID NO: 2; is a natural polymorphic variant of SEQ ID NO: 2; has a length at least about 30 amino acids; exhibits at least two non-overlapping epitopes which are specific for a primate DCRS5; has a molecular weight of at least 30 kD with natural glycosylation; is a synthetic polypeptide; is in a sterile form; is in an aqueous or buffered solution; is attached to a solid substrate; is conjugated to another chemical moiety; or is physically associated with an  
20   IL-12R $\beta$ 1 polypeptide.

Other embodiments of the invention provide: a substantially pure or recombinant polypeptide comprising at least two distinct nonoverlapping segments of at least six contiguous amino acids of the intracellular portion of SEQ ID NO: 2; a substantially pure or recombinant polypeptide comprising at least twelve contiguous amino acids of the intracellular portion of  
25   SEQ ID NO: 2; or a substantially pure natural sequence polypeptide comprising mature SEQ ID NO: 2. In particular forms, the polypeptide comprising at least two distinct nonoverlapping segments of at least six contiguous amino acids of the intracellular portion of SEQ ID NO: 2 will be where: the distinct nonoverlapping segments: include one of at least twelve amino acids; include one of at least seven amino acids and a second of at least nine amino acids;  
30   include a third distinct segment of at least six amino acids; or comprise one of R355-L373, P378-L405, V407-D426, K428-D439, P441-V452, I454-G460, I465-T587, or N592-606; or the polypeptide further comprises at least two distinct nonoverlapping segments of at least six contiguous amino acids of the extracellular portion of SEQ ID NO: 2. Alternatively, the polypeptide comprising at least twelve contiguous amino acids of the intracellular portion of  
35   SEQ ID NO: 2 will be one where: the at least twelve contiguous amino acid segment comprises one of R355-L373, P378-L405, V407-D426, K428-D439, P441-V452, I454-G460, I465-T587,

or N592-606; or the polypeptide further comprises at least two distinct nonoverlapping segments of at least six contiguous amino acids of the extracellular portion of SEQ ID NO: 2. Or, the pure natural sequence polypeptide comprising mature SEQ ID NO: 2 may further comprising a purification or detection epitope. Such polypeptides may: consist of the mature  
5 sequence of Table 1; be an unglycosylated polypeptide; be from a human; comprise at least 40 contiguous amino acids of SEQ ID NO: 2; exhibit at least three nonoverlapping segments of at least fifteen contiguous amino acids of SEQ ID NO: 2; be a natural polymorphic variant of SEQ ID NO: 2; have a length at least about 30 amino acids; exhibit at least two non-overlapping epitopes which are specific for a primate DCRS5; have a molecular weight of at  
10 least 30 kD with natural glycosylation; be a synthetic polypeptide; be in a steril form; be in an aqueous or buffered solution; be attached to a solid substrate; be conjugated to another chemical moiety; or be physically associated with an IL-12R $\beta$ 1 polypeptide.

Various other compositions are provided, e.g., comprising: a substantially pure polypeptide combined with the IL-12R $\beta$ 1 protein; or such a polypeptide in a carrier, wherein  
15 the carrier is: an aqueous compound, including water, saline, and/or buffer; and/or formulated for oral, rectal, nasal, topical, or parenteral administration.

Kits are provided comprising such a polypeptide and: a compartment comprising the polypeptide; a compartment comprising an IL-12R $\beta$ 1 polypeptide; a compartment comprising a p40, IL-B30, or p40/IL-B30 polypeptide; or instructions for use or disposal of reagents in the  
20 kit.

Antibodies and other binding compounds are provided, e.g., comprising an antigen binding site from an antibody, which specifically binds to the intracellular portion of the DCRS5, wherein: the binding compound is in a container; the polypeptide is from a human; the binding compound is an Fv, Fab, or Fab2 fragment; the binding compound is conjugated to  
25 another chemical moiety; or the antibody: is raised against a peptide sequence of a mature polypeptide of Table 1; is raised against a mature DCRS5; is raised to a purified human DCRS5; is immunoselected; is a polyclonal antibody; binds to a denatured DCRS5; exhibits a Kd to antigen of at least 30  $\mu$ m; is attached to a solid substrate, including a bead or plastic membrane; is in a sterile composition; or is detectably labeled, including a radioactive or  
30 fluorescent label. Kits are also provided comprising the binding compound and: a compartment comprising the binding compound; a compartment comprising: a p40 polypeptide; an IL-B30 polypeptide; a DCRS5 polypeptide; and/or an IL-12R $\beta$ 1 polypeptide; a compartment comprising an antibody which binds selectively to: a p40 polypeptide; an IL-B30 polypeptide; a DCRS5 polypeptide; and/or an IL-12R $\beta$ 1 polypeptide; or instructions for  
35 use or disposal of reagents in the kit.

Also provided are methods, e.g., of producing an antigen:antibody complex, comprising contacting under appropriate conditions a primate DCRS5 polypeptide with an antibody, thereby allowing the complex to form. Such method may be where: the complex is purified from other cytokine receptors; the complex is purified from other antibody; the  
5 contacting is with a sample comprising an interferon; the contacting allows quantitative detection of the antigen; the contacting is with a sample comprising the antibody; or the contacting allows quantitative detection of the antibody. Other compositions are provided, e.g., composition comprising: a sterile binding compound, or the binding compound and a carrier, wherein the carrier is: an aqueous compound, including water, saline, and/or buffer;  
10 and/or formulated for oral, rectal, nasal, topical, or parenteral administration.

The invention also provides an isolated or recombinant nucleic acid encoding the DCRS5 polypeptide, wherein the: DCRS5 is from a human; or the nucleic acid: encodes an antigenic peptide sequence of Table 1; encodes a plurality of antigenic peptide sequences of Table 1; exhibits identity over at least thirteen nucleotides to a natural cDNA encoding the  
15 segment; is an expression vector; further comprises an origin of replication; is from a natural source; comprises a detectable label; comprises synthetic nucleotide sequence; is less than 6 kb, preferably less than 3 kb; is from a primate; comprises a natural full length coding sequence; is a hybridization probe for a gene encoding the DCRS5; or is a PCR primer, PCR product, or mutagenesis primer. Cells comprising the recombinant nucleic acid are provided,  
20 including where the cell is: a prokaryotic cell; a eukaryotic cell; a bacterial cell; a yeast cell; an insect cell; a mammalian cell; a mouse cell; a primate cell; or a human cell.

Kit embodiments include those comprising the nucleic acid and: a compartment comprising the nucleic acid; a compartment comprising a nucleic acid encoding: a p40 polypeptide; an IL-B30 polypeptide; a DCRS5 polypeptide; and/or an IL-12R $\beta$ 1 polypeptide; a  
25 compartment comprising: a p40 polypeptide; an IL-B30 polypeptide; a DCRS5 polypeptide; and/or an IL-12R $\beta$ 1 polypeptide; a compartment comprising an antibody which selectively binds to: a p40 polypeptide; an IL-B30 polypeptide; a DCRS5 polypeptide; and/or an IL-12R $\beta$ 1 polypeptide; or instructions for use or disposal of reagents in the kit.

Other nucleic acid embodiments include those which: hybridize under wash conditions  
30 of 30 minutes at 30° C and less than 2M salt to the portion of SEQ ID NO: 1 encoding the intracellular portion; or exhibit identity over a stretch of at least about 30 nucleotides to the intracellular portion of a primate DCRS5. Preferably, such nucleic acid will be one wherein: the wash conditions are at 45° C and/or 500 mM salt; or 55° C and/or 150 mM salt; or the stretch is at least 55 or 75 nucleotides.

35 Therapeutic uses include methods of modulating physiology or development of a cell comprising contacting the cell with: an antagonist of p40/IL-B30 which is a complex

comprising: the extracellular portion of a primate DCRS5 and/or the extracellular portion of a primate IL-12R $\beta$ 1; an antagonist of p40/IL-B30 which is an antibody which binds a complex comprising: primate DCRS5 and/or primate IL-12R $\beta$ 1; an antagonist of p40/IL-B30 which is an antibody which binds to DCRS5; an antagonist of p40/IL-B30 which is an antibody to IL-12R $\beta$ 1; an antagonist of p40/IL-B30 which is an antisense nucleic acid to DCRS5 or IL-12R $\beta$ 1; or an agonist of p40/IL-B30 which is an antibody which binds a complex comprising primate DCRS5 and/or primate IL-12R $\beta$ 1. In one type of method, the contacting is with an antagonist, and the contacting is in combination with an antagonist to IL-12, IL-18, TNF, and/or IFN $\gamma$ ; or the cell is from a host which: exhibits signs or symptoms of a chronic TH1 mediated disease; exhibits symptoms or signs of multiple sclerosis, rheumatoid arthritis, osteoarthritis, inflammatory bowel disease, diabetes, psoriasis, or sepsis; or receives an allogeneic transplant. Conversely, the method may be contacting with an agonist, and: the contacting is in combination with IL-12, IL-18, TNF, or IFN $\gamma$ ; or the cell is from a host which: exhibits signs or symptoms of a chronic Th2 response; suffers from a tumor, viral, or fungal growth; receives a vaccine; or suffers from an allergic response.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

### OUTLINE

- I. General
- II. Activities
- III. Nucleic acids
  - A. encoding fragments, sequence, probes
  - B. mutations, chimeras, fusions
  - C. making nucleic acids
  - D. vectors, cells comprising
- IV. Proteins, Peptides
  - A. fragments, sequence, immunogens, antigens
  - B. muteins
  - C. agonists/antagonists, functional equivalents
  - D. making proteins
- V. Making nucleic acids, proteins
  - A. synthetic
  - B. recombinant
  - C. natural sources
- VI. Antibodies
  - A. polyclonals
  - B. monoclonal
  - C. fragments; Kd
  - D. anti-idiotypic antibodies
  - E. hybridoma cell lines

VII. Kits, Diagnosis, and Quantitation

- A. ELISA
- B. assay mRNA encoding
- 5 C. qualitative/quantitative
- D. kits

VIII. Therapeutic compositions, methods

- A. combination compositions
- B. unit dose
- 10 C. administration

IX. Screening

I. General

20 The present invention provides the amino acid sequence and DNA sequence of mammalian, herein primate, cytokine receptor-like subunit molecules, this one designated DNAX Cytokine Receptor Subunit 5 (DCRS5) having particular defined properties, both structural and biological. Various cDNAs encoding these molecules were obtained from primate, e.g., human, cDNA sequence libraries. Other primate or other mammalian counterparts would also be desired.

20 Additionally, the invention provides matching of the p40/IL-B30 ligand with receptor subunits DCRS5 and IL-12R $\beta$ 1, which pairing provides insight into indications for use of the agonists and antagonists based upon reagents directed thereto.

25 Some of the standard methods applicable are described or referenced, e.g., in Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols. 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and periodic supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York; each of which is incorporated herein by reference.

30 Nucleotide (SEQ ID NO: 1) and corresponding amino acid sequence (SEQ ID NO: 2) of a primate, e.g., human, DCRS5 coding segment is shown in Table 1. The predicted signal sequence is indicated, but may depend on cell type, or may be a few residues in either direction. Potential N glycosylation sites are at Asparagine residues 6, 24, 58, 118, 157, 209, and 250. Disulfide linkages are likely to be found between cysteine residues at positions 29 and 78; and a conserved C<sub>2</sub>CXW motif is found at positions 110/121/123. The tryptophan at 35 219; and the WxxWS motif from 281-285 are notable. The segment from about 1-101 is an Ig domain; from about 102-195 is a cytokine binding domain 1; from about 196-297 is a cytokine binding domain 2; from about 298-330 is a linker; from about 329-354 is a transmembrane segment; and from about 356-606 is an intracellular domain. Intracellular features include



putative SH2 binding sites at Y374-I377, Y461-Q464, and Y588-Q591; and potentially important tyrosine residues at 406, 427, 440, and 453. These sites and boundaries are notable.

The ORF contains a putative signal sequence which is predicted to be cleaved at ...CHG/GIT... as shown above. A predicted extracellular domain of 328 amino acids is  
5 followed by a putative transmembrane segment, and finally a cytoplasmic domain of about 252 amino acids. The ligand-binding functions are predicted to reside in the extracellular domain.

The reverse translation nucleic acid sequence is provided in Table 2.

Table 1: Nucleotide and polypeptide sequences of DNAX Cytokine Receptor Subunit like embodiments (DCRS5). Primate, e.g., human embodiment (see SEQ ID NO: 1 and 2). Predicted signal sequence indicated, but may vary by a few positions and depending upon cell type. Identified positions of variation are at nucleotides 127 and 563; which are paired G and G (translating to the combination of Q and G) or T and A (translating to H and R).

5	gtggtacggg aattccattg tgttgggcag ccaacaaggg tggcagcctg gctctgaagt 60	
10	ggaattatgt gcttcaaaca gggtgaaaga gggaaacagt cttttcctgc ttccagac 118	
	atg aat cak gtc act att caa tgg gat gca gta ata gcc ctt tac ata 166	
	Met Asn Xaa Val Thr Ile Gln Trp Asp Ala Val Ile Ala Leu Tyr Ile	
	-20 -15 -10	
15	ctc ttc agc tgg tgt cat gga gga att aca aat ata aac tgc tct ggc 214	
	Leu Phe Ser Trp Cys His Gly Gly Ile Thr Asn Ile Asn Cys Ser Gly	
	-5 -1 1 5	
20	cac atc tgg gta gaa cca gcc aca att ttt aag atg ggt atg aat atc 262	
	His Ile Trp Val Glu Pro Ala Thr Ile Phe Lys Met Gly Met Asn Ile	
	10 15 20 25	
25	tct ata tat tgc caa gca gca att aag aac tgc caa cca agg aaa ctt 310	
	Ser Ile Tyr Cys Gln Ala Ala Ile Lys Asn Cys Gln Pro Arg Lys Leu	
	30 35 40	
30	cat ttt tat aaa aat ggc atc aaa gaa aga ttt caa atc aca agg att 358	
	His Phe Tyr Lys Asn Gly Ile Lys Glu Arg Phe Gln Ile Thr Arg Ile	
	45 50 55	
	aat aaa aca aca gct cgg ctt tgg tat aaa aac ttt ctg gaa cca cat 406	
	Asn Lys Thr Thr Ala Arg Leu Trp Tyr Lys Asn Phe Leu Glu Pro His	
	60 65 70	
35	gct tct atg tac tgc act gct gaa tgt ccc aaa cat ttt caa gag aca 454	
	Ala Ser Met Tyr Cys Thr Ala Glu Cys Pro Lys His Phe Gln Glu Thr	
	75 80 85	
40	ctg ata tgt gga aaa gac att tct tct gga tat ccg cca gat att cct 502	
	Leu Ile Cys Gly Lys Asp Ile Ser Ser Gly Tyr Pro Pro Asp Ile Pro	
	90 95 100 105	
45	gat gaa gta acc tgt gtc att tat gaa tat tca ggc aac atg act tgc 550	
	Asp Glu Val Thr Cys Val Ile Tyr Glu Tyr Ser Gly Asn Met Thr Cys	
	110 115 120	
50	acc tgg aat gct rgg aag ctc acc tac ata gac aca aaa tac gtg gta 598	
	Thr Trp Asn Ala Xaa Lys Leu Thr Tyr Ile Asp Thr Lys Tyr Val Val	
	125 130 135	
	cat gtg aag agt tta gag aca gaa gaa gag caa cag tat ctc acc tca 646	
	His Val Lys Ser Leu Glu Thr Glu Glu Glu Gln Gln Tyr Leu Thr Ser	
	140 145 150	
55		

	agc	tat	att	aac	atc	tcc	act	gat	tca	tta	caa	ggc	ggc	aag	aag	tac	694
	Ser	Tyr	Ile	Asn	Ile	Ser	Thr	Asp	Ser	Leu	Gln	Gly	Gly	Lys	Lys	Tyr	
	155						160					165					
5	ttg	gtt	tgg	gtc	caa	gca	gca	aac	gca	cta	ggc	atg	gaa	gag	tca	aaa	742
	Leu	Val	Trp	Val	Gln	Ala	Ala	Asn	Ala	Leu	Gly	Met	Glu	Glu	Ser	Lys	
	170					175					180					185	
10	caa	ctg	caa	att	cac	ctg	gat	gat	ata	gtg	ata	cct	tct	gca	gcc	gtc	790
	Gln	Leu	Gln	Ile	His	Leu	Asp	Asp	Ile	Val	Ile	Pro	Ser	Ala	Ala	Val	
					190					195					200		
15	att	tcc	agg	gct	gag	act	ata	aat	gct	aca	gtg	ccc	aag	acc	ata	att	838
	Ile	Ser	Arg	Ala	Glu	Thr	Ile	Asn	Ala	Thr	Val	Pro	Lys	Thr	Ile	Ile	
				205					210					215			
20	tat	tgg	gat	agt	caa	aca	aca	att	gaa	aag	gtt	tcc	tgt	gaa	atg	aga	886
	Tyr	Trp	Asp	Ser	Gln	Thr	Thr	Ile	Glu	Lys	Val	Ser	Cys	Glu	Met	Arg	
			220					225					230				
25	tac	aag	gct	aca	aca	aac	caa	act	tgg	aat	gtt	aaa	gaa	ttt	gac	acc	934
	Tyr	Lys	Ala	Thr	Thr	Asn	Gln	Thr	Trp	Asn	Val	Lys	Glu	Phe	Asp	Thr	
		235				240					245						
30	aat	ttt	aca	tat	gtg	caa	cag	tca	gaa	ttc	tac	ttg	gag	cca	aac	att	982
	Asn	Phe	Thr	Tyr	Val	Gln	Gln	Ser	Glu	Phe	Tyr	Leu	Glu	Pro	Asn	Ile	
	250					255					260					265	
35	aag	tac	gta	ttt	caa	gtg	aga	tgt	caa	gaa	aca	ggc	aaa	agg	tac	tgg	1030
	Lys	Tyr	Val	Phe	Gln	Val	Arg	Cys	Gln	Glu	Thr	Gly	Lys	Arg	Tyr	Trp	
				270					275						280		
40	cag	cct	tgg	agt	tca	ccg	ttt	ttt	cat	aaa	aca	cct	gaa	aca	gtt	ccc	1078
	Gln	Pro	Trp	Ser	Ser	Pro	Phe	Phe	His	Lys	Thr	Pro	Glu	Thr	Val	Pro	
				285					290					295			
45	cag	gtc	aca	tca	aaa	gca	ttc	caa	cat	gac	aca	tgg	aat	tct	ggg	cta	1126
	Gln	Val	Thr	Ser	Lys	Ala	Phe	Gln	His	Asp	Thr	Trp	Asn	Ser	Gly	Leu	
			300					305					310				
50	aca	gtt	gct	tcc	atc	tct	aca	ggg	cac	ctt	act	tct	gac	aac	aga	gga	1174
	Thr	Val	Ala	Ser	Ile	Ser	Thr	Gly	His	Leu	Thr	Ser	Asp	Asn	Arg	Gly	
		315					320					325					
55	gac	att	gga	ctt	tta	ttg	gga	atg	atc	gtc	ttt	gct	gtt	atg	ttg	tca	1222
	Asp	Ile	Gly	Leu	Leu	Leu	Gly	Met	Ile	Val	Phe	Ala	Val	Met	Leu	Ser	
	330					335				340					345		
60	att	ctt	tct	ttg	att	ggg	ata	ttt	aac	aga	tca	ttc	cga	act	ggg	att	1270
	Ile	Leu	Ser	Leu	Ile	Gly	Ile	Phe	Asn	Arg	Ser	Phe	Arg	Thr	Gly	Ile	
				350					355						360		
65	aaa	aga	agg	atc	tta	ttg	tta	ata	cca	aag	tgg	ctt	tat	gaa	gat	att	1318
	Lys	Arg	Arg	Ile	Leu	Leu	Leu	Ile	Pro	Lys	Trp	Leu	Tyr	Glu	Asp	Ile	
				365					370					375			

	cct aat atg aaa aac agc aat gtt gtg aaa atg cta cag gaa aat agt	1366
	Pro Asn Met Lys Asn Ser Asn Val Val Lys Met Leu Gln Glu Asn Ser	
	380 385 390	
5	gaa ctt atg aat aat aat tcc agt gag cag gtc cta tat gtt gat ccc	1414
	Glu Leu Met Asn Asn Asn Ser Ser Glu Gln Val Leu Tyr Val Asp Pro	
	395 400 405	
10	atg att aca gag ata aaa gaa atc ttc atc cca gaa cac aag cct aca	1462
	Met Ile Thr Glu Ile Lys Glu Ile Phe Ile Pro Glu His Lys Pro Thr	
	410 415 420 425	
15	gac tac aag aag gag aat aca gga ccc ctg gag aca aga gac tac ccg	1510
	Asp Tyr Lys Lys Glu Asn Thr Gly Pro Leu Glu Thr Arg Asp Tyr Pro	
	430 435 440	
20	caa aac tcg cta ttc gac aat act aca gtt gta tat att cct gat ctc	1558
	Gln Asn Ser Leu Phe Asp Asn Thr Thr Val Val Tyr Ile Pro Asp Leu	
	445 450 455	
25	aac act gga tat aaa ccc caa att tca aat ttt ctg cct gag gga agc	1606
	Asn Thr Gly Tyr Lys Pro Gln Ile Ser Asn Phe Leu Pro Glu Gly Ser	
	460 465 470	
30	cat ctc agc aat aat aat gaa att act tcc tta aca ctt aaa cca cca	1654
	His Leu Ser Asn Asn Asn Glu Ile Thr Ser Leu Thr Leu Lys Pro Pro	
	475 480 485	
35	gtt gat tcc tta gac tca gga aat aat ccc agg tta caa aag cat cct	1702
	Val Asp Ser Leu Asp Ser Gly Asn Asn Pro Arg Leu Gln Lys His Pro	
	490 495 500 505	
40	aat ttt gct ttt tct gtt tca agt gtg aat tca cta agc aac aca ata	1750
	Asn Phe Ala Phe Ser Val Ser Ser Val Asn Ser Leu Ser Asn Thr Ile	
	510 515 520	
45	ttt ctt gga gaa tta agc ctc ata tta aat caa gga gaa tgc agt tct	1798
	Phe Leu Gly Glu Leu Ser Leu Ile Leu Asn Gln Gly Glu Cys Ser Ser	
	525 530 535	
50	cct gac ata caa aac tca gta gag gag gaa acc acc atg ctt ttg gaa	1846
	Pro Asp Ile Gln Asn Ser Val Glu Glu Glu Thr Thr Met Leu Leu Glu	
	540 545 550	
55	aat gat tca ccc agt gaa act att cca gaa cag acc ctg ctt cct gat	1894
	Asn Asp Ser Pro Ser Glu Thr Ile Pro Glu Gln Thr Leu Leu Pro Asp	
	555 560 565	
60	gaa ttt gtc tcc tgt ttg ggg atc gtg aat gag gag ttg cca tct att	1942
	Glu Phe Val Ser Cys Leu Gly Ile Val Asn Glu Glu Leu Pro Ser Ile	
	570 575 580 585	
65	aat act tat ttt cca caa aat att ttg gaa agc cac ttc aat agg att	1990
	Asn Thr Tyr Phe Pro Gln Asn Ile Leu Glu Ser His Phe Asn Arg Ile	
	590 595 600	

tca ctc ttg gaa aag tagagctgtg tgggtcaaaat caatatgaga aagctgcctt 2045  
Ser Leu Leu Glu Lys  
605

5 gcaatctgaa cttgggtttt ccctgcaata gaaattgaat tctgcctctt tttgaaaaaa 2105  
atgtattcac atacaaatct tcacatggac acatgttttc atttccttg gataaatacc 2165  
taggtagggg attgctgggc catatgataa gcatatgttt cagttctacc aatcttgttt 2225  
10 ccagagtagt gacatttctg tgctcctacc atcaccatgt aagaattccc gggagctcca 2285  
tgccttttta attttagcca ttcttctgcc tmatttctta aaattagaga attaaggtcc 2345  
15 cgaaggtgga acatgcttca tggtcacaca tacaggcaca aaaacagcat tatgtggacg 2405  
cctcatgtat tttttataga gtcaactatt tcctctttat tttccctcat tgaaagatgc 2465  
aaaacagctc totattgtgt acagaaaggg taaataatgc aaaataacctg gtagtaaaat 2525  
20 aaatgctgaa aattttcctt taaaatagaa tcattaggcc aggcgtgggtg gctcatgctt 2585  
gtaatcccag cactttggta ggctgaggtr ggtggatcac ctgaggtcag gagttcgagt 2645  
25 ccagcctggc caatatgctg aaaccctgtc tctactaaaa ttacaaaaat tagccggcca 2705  
tgggtggcagg tgcttgtaat ccagctact tgggaggctg aggcaggaga atcacttgaa 2765  
ccaggaaggc agaggttgca ctgagctgag attgtgccac tgcactccag cctgggcaac 2825  
30 aagagcaaaa ctctgtctgg aaaaaaaaaa aaaa 2859

35 MN (O/H) VTIOWDVIALYILFSWCHGGITNINCSGHIWVEPATIFKMGMNISIYCQAAIKNCQPRKLHF  
YKNGIKERFQITRINKTTARLWYKNFLEPHASMYCTAECPKHFQETLICGKDISSGYPPDIPDEVTCVIY  
EYSGNMTCTWNA (G/R) KLTYIDTKYVVHVKSLETEEEQQYLTSSYINISTDSLQGGKKYLVWVQAANAL  
GMEESKQLQIHLDDIVIPSAAVISRAETINATVPKTIYWDQSQTIEKVSCEMRYKATTNQTWNVKEFDT  
NFTYVQQSEFYLEPNIKYVFQVRCQETGKRYWQPWSSPFFHKTPETVPQVTSKAFQHDTWNSGLTVASIS  
TGHLTSDNRGDIGLLGMIVFAVMLSILSLIGIFNRSFRTGIKRRILLIPKWLYEDIPNMKNSNVVKML  
40 QENSELMNNSSEQVLYVDPMITEIKEIFIPEHKPTDYKENTGPLETRDYPQNSLFDNTTVVYIPDLNT  
GYKPQISNFLPEGSHLSNNNEITSLTLKPPVDSLDSGNNPRLQKHPNFAFSVSSVNSLSNTIFLGELSLI  
LNQGECSSPDIQNSVEEETTMLLENDSPSETIPEQTLLPDEFVSLGIVNEELPSINTYFPQNILESHFN  
RISLLEK

Table 2: Reverse Translation of primate, e.g., human, DCRS5 (SEQ ID NO: 3):

ATGAAYCAYGTNACNATHCARTGGGAYGCNGTNATHGCNYTNTAYATHYTNNTTYWSNTGGTGYCAYGGNGGNAT  
 5 HACNAAYATHAAYTGYWSNGGNCAYATHTGGGTNGARCCNGCNACNATHHTTYAARATGGGNATGAAYATHWSNA  
 THTAYTGYCARGCNGCNATHAARAAYTGYCARCCNMGNAARYTNCAYTTYTAYAARAAYGGNATHAARGARMGN  
 TTYCARATHACNMGNATHAAYAARACNACNGCNMGNYTNTGGTAYAARAAYTTYTNGARCCNCAYGCNWSNAT  
 GTAYTGYACNGCNGARTGYCCNAARCAYTTYCARGARACNYTNATHGTGYGGNAARGAYATHWSNWSNGGNTAYC  
 CNCCNGAYATHCCNGAYGARGTNACNTGYGTNATHTAYGARTAYWSNGGNAAYATGACNTGYACNTGGAAYGCN  
 10 MGNAARYTNACNTAYATHGAYACNAARTAYGTNGTNCAYGTNAARWSNYTNGARACNGARGARGARCARCARTA  
 YYTNACNWSNWSNTAYATHAAYATHWSNACNGAYWSNYTNCARGGNGGNAARAARTAYYTNGTNTGGGTNCARG  
 CNGCNAAYGCNYTNGGNATGGARGARWSNAARCARYTNCARATHCAYYTNGAYGAYATHGTNATHCCNWSNGCN  
 GCNGTNATHWSNMGNGCNGARACNATHAAYGCNACNGTNCNAARACNATHATHTAYTGGGAYWSNCARACNAC  
 NATHGARAARGTNWSNTGYGARATGMGNTAYAARGCNACNACNAAYCARACNTGGAAYGTNAARGARTTYGAYA  
 CNAAYTTYACNTAYGTNCARCARWSNGARTTYTAYYTNGARCCNAAYATHAARTAYGTNTTYCARGTNMGNTGY  
 15 CARGARACNGGNAARMGNTAYTGGCARCCTGGWSNWSNCCNTTYTTYCAYAAACNCCNGARACNGTNCNCNA  
 RGTNACNWSNAARGCNTTYCARCAYGAYACNTGGAAYWSNGGNYTNACNGTNGCNWSNATHWSNACNGGNCAYY  
 TNACNWSNGAYAAVMGNGGNGAYATHGGNYTNYTNYTNGGNATGATHGTNTTYGCNGTNATGYTNWSNATHYTN  
 WSNYTNATHGGNATHHTTYAAYMGNWSNTTYMGNACNGGNATHAARMGMNMGNATHYTNNTNYTNATHCCNAARTG  
 20 GYTNTAYGARGAYATHCCNAAYATGAARAAYWSNAAYGTNGTNAARATGYTNCARGARAAYWSNGARYTNATGA  
 AYAAYAAYWSNWSNGARCARGTNYTNTAYGTNGAYCCNATGATHACNGARATHAARGARATHHTTYATHCCNGAR  
 CAYAARCCNACNGAYTAYAARAARGARAAYACNGGNCNYTNGARACNMGNGAYTAYCCNCARAAYWSNYTNTT  
 YGAYAAACNACNGTNGTNTAYATHCCNGAYYTNAAYACNGGNTAYAARCCNCARATHWSNAAYTTYYTNCNG  
 ARGGNWSNCAYYTNWSNAAYAAAYAGARATHACNWSNYTNACNYTNAARCCNCCNGTNGAYWSNYTNGAYWSN  
 25 GGNAAYAAACNMGNYTNCARAARCAAYCCNAAYTTYGCNTTYWSNGTWSNWSNGTNAAYWSNYTNWSNAAYAC  
 NATHHTTYTNGGNGARYTNWSNYTNATHYTNAAYCARGGNGARTGYWSNWSNCCNGAYATHCARAAYWSNGTNG  
 ARGARGARACNACNATGYTNYTNGARAAYGAYWSNCCNWSNGARACNATHCCNGARACARACNYTNYTNCNGAY  
 GARTTYGTNWSNTGYTNGGNATHGTNAAYGARGARYTNCCNWSNATHAAYACNTAYTTYCCNCARAAYATHYT  
 NGARWSNCAYYTYAAYMGNATHWSNYTNYTNGARAAR

30

Table 3: Alignment of various cytokine receptor subunits. Human IL-6 receptor protein gp130 is SEQ ID NO: 4 (GenBank M57230); human IL-12 receptor beta2 subunit is SEQ ID NO: 5 (GenBank U64198).

35	huIL-12R 2	1	MAHTFRGCSLAFMFIITWLLIKAKIDACKRGDVTVPKPSHVILLGSTVN	48
	hugp130	1	MLTLQTWVQALFIFLTTESTGELLDPG---YISPESPVVQLHSNFT	45
	huDCRS5	1	MNHVTIQWDAVIALYILFSWCHGGITNINCS-GHIWVEPATIFKMGMNIS	49
			* * * * *	
40	huIL-12R 2	49	ITCSLKPRQGCIFYSRNRKLILYKFDRRINFHHGHSLSNQVTGLPLG---	95
	hugp130	46	AVCVLKEKCMDYFHVNANYIVWKTNHFTIPKEQYTIINRTASSVTFTDIA	95
	huDCRS5	50	IYCQAAIKN--CQP---RKLHFKNGIKER-FQITRINKTTARLWYKNFL	93
			* * * * *	
45	huIL-12R 2	96	--TTLFVCKLACINSD-BIQICGAEIFVGVAPQPNLSCIQKEQGTVA	142
	hugp130	96	SLNIQLTCNLTFGQL-EQNVYGITIISGLPPEKPKNLSCIVN-EGKKMR	143
	huDCRS5	94	EPHASYCTAECPKHFQETLICGKDISSGYPPDIPDEVTCVIYEYSNMT	143
			* * * * *	
50	huIL-12R 2	143	CTWERGRDTHLYTEYTLQLSGPKNLTWQKQCKDIYCDYLDGFINLTPESP	192
	hugp130	144	CEWDGGRETHLETNFTLKS--EWATHKFADCKAKRDTPTSCTVDYS-TVY	190
	huDCRS5	144	CTWNARKLTYIDTKYVVHVKSLETEEQYLTSSYINISTDSLQGG----	189
			* * * * *	
55	huIL-12R 2	193	ESNFTAKVTAVNSLGSSSSLPSTFTFLDIVRPLPPWDIRIKFQKASVSR	242
	hugp130	191	FVNIEVWVEAENALGKVTSDFINFDVPYKVKPNPPHNLSVINSEELSSIL	240

	huDCRS5	190	-KKYLVWVQAANALGMEESKQLQIHLDDIVIPSAAVISRAETINATVPKT	238
			* * * . * * *	
5	huIL-12R	2 243	TLYWRD----EGLVLLNRLRYRPSNSRLWMNVN---VTKAKGRHDLDDLK	285
	hugp130	241	KLWTNPSIKSVIILKYNIQYRTKDASTWSQIPPEDTASTRSSFTVQDLK	290
	huDCRS5	239	IIYWDS--QTTIEKVSCEMRYKATTNQTWNVKEFD-TNFTYVQQSEFYLE	285
			* . . . . *	
10	huIL-12R	2 286	PFTEYEFQISSKLHLYKGSWSDWSESLRAQTPEEEPTGMLDVWYMKRHID	335
	hugp130	291	PFTEYVFRIRCMKEDGKGYSDWSEEASGITYEDRPSKAPSFYWKIDPSH	340
	huDCRS5	286	PNIKYVFQVRCQ-ETGKRYWQPWSSPFFHKTPETVP-----	320
			* * * . . * * *	
15	huIL-12R	2 336	YS-RQQISLFWKNLSVSEARGKILHYQVTLQELTGGKAMTONITGHTSWT	384
	hugp130	341	TQGYRTVQLVWKTLPPEFANGKILDYEV---LTRWKSHLQNYTVNATKL	387
	huDCRS5	321	-----QVTSKAFQHDTWNSGLTVASISTG-----HLTSDN--RGDIGLL	357
			. . . . *	
20	huIL-12R	2 385	TVIPRTGNWAVAVSAANSKGSSLPTRINIMNLCEAGLLAPROVSANSEGM	434
	hugp130	388	TVNLTNDRYLATLTVRNLVGKSDAAVLTIP-ACDFQATHPVMDLKAFPDK	436
	huDCRS5	358	LGMIVFAVMLSILSLIGIFNRSFRTGIKRR-----	387
			. . . . *	
25	huIL-12R	2 435	DNILVTWQPPRKDPSAVQYVVEWRELHPG-GDTQVPLNWLRSRPNVSA	483
	hugp130	437	NMLWVEWTTTPRE---SVKKYILEWCVLS---DKAPCITDWQQEDGTVHRT	480
	huDCRS5	388	-----ILLIPKWLYEDIPNMKNSNVVKMLQEN---SE	417
			. . . . *	
30	huIL-12R	2 484	LISENIKSYICYEIRVYALSGDQ-GGCSSILGNSKHKAPLSGPHINAITE	532
	hugp130	481	YLRGNLAESKCYLITVTPVYADGPGSPESIKAYLKQAPPSKGPTVRTKKV	530
	huDCRS5	418	LMNNNSSE-----QVLYVDP-----MITEIKEIFIPEHKPTDYKKE-	453
			* . . . . *	
35	huIL-12R	2 533	EKGSILISWNSIPVQEQMGCLLHYRIYWKERDSNSQPQLCEIPYRVSQNS	582
	hugp130	531	GKNEAVLEWDQLPVDVQNGFIRNYTIFYRTIIGN---ETAVNVDSSTE	576
	huDCRS5	454	--NTGPLETRDYP---QNSLFDNTTVVYIPDLNTG-----YKPQISN--	490
			* * . . . *	
40	huIL-12R	2 583	HPINSLQPRVTVYVLWMTALTAAGESSHGNEREFCLQGGKAN-WMAFVAPSI	631
	hugp130	577	YTLSSLTSDTLYMVRMAAYTDEG-GKDGPEFTFTTPKFAQGEIEAIVPV	625
	huDCRS5	491	-----FLPEG-----	495
			. . . . *	
45	huIL-12R	2 632	CIAIIMVGIFSTHYFQQKFVLLAALRP-----QWCSREIPDPA	670
	hugp130	626	CLAFLLTTLGVLFCFNKRDLIKKHIWPNVPDPSKSHIAQWSPHTPPRH	675
	huDCRS5	496	-----SHLSNNN-EITSLTLKP-----PVDSLDG	519
			. . . . *	
50	huIL-12R	2 671	NSTCAKKYPIAEETQLPLDRLLID-WPTPEDPEPLVIS--EVLHQVTPV	717
	hugp130	676	FNSKDQMYSDGNFTDVSVEIEANDKKPFPEDLKSLDLFKKEKINTEGHS	725
	huDCRS5	520	NNPRLQKHFN-FAFSVSSVNSLSNT-----I---FLGELSLI	552
			. . . . *	
55	huIL-12R	2 718	FRHPPCSNWPQREKGIQGHQASEKDMMHSASSPPPPRALQAESRQLVDLY	767
	hugp130	726	SGIGSSCMSSSRPSISSSDENESSQNTSSTVQYSTVVHSGYRHQVPSVQ	775
	huDCRS5	553	LNQGECS---S---PDIQNSVEEETMLLENDSP-----	580
			* . . . . *	

5	huIL-12R	2	768	KVLESRGSDPKPENPACPWTVLPAGDLPTHGGLPSN---IDDLPSHEAP	814
	hugp130		776	VFSRSESTQPLLDSEERPDLQLVDHVDGGDGLPRQQYFKQNCSEQHES	825
	huDCRS5		581	--SETIPEQTLLPDEFVSLGIVNEELPSINTYFPQN---ILESHEFNR--	623
				*	
10	huIL-12R	2	815	LADSLEEELEPQHISLS-----VFPSSSLHPLTFSCG-----	845
	hugp130		826	PDISHFERSKQVSSVNEEDFVRLKQQISDHISQSCGSGQMMPQEVSAAD	875
	huDCRS5		624	--ISLLEK	629
				* *	
15	huIL-12R	2	846	-----DKLTLDQLKMRCDSLML	862
	hugp130		876	AFGPGTEGQVERFETVGMEEAATDEGMPKSYLPQTVRQGGYMPQ	918
	huDCRS5		630		629

The closest relatives of the extracellular domain of "IL-30R" are the IL-6 signal transducer gp130 and IL-12R $\beta$ 2. Somewhat less close relatives are GCSF receptor, leptin receptor, leukemia inhibitory factor receptor, and CNTF receptor. Thus "IL-30R" is a member of the class I branch of the cytokine receptor superfamily and is closely related to the IL-6R/IL-12R family.

Table 3 shows comparison of the available sequences of primate receptor subunits with the primate, e.g., human DCRS5 (IL-30R). The DCRS5 shows similarity to the IL-6 receptor subunit gp130 (e.g., IL-6R subunit) and the IL-12R $\beta$ 2 subunit. The DCRS5 exhibits structural features of a beta subunit, but the actual sequence of protein interactions and signaling remains unresolved.

As used herein, the term DCRS5 shall be used to describe a protein comprising the amino acid sequence shown in Table 1. In many cases, a substantial fragment thereof will be functionally or structurally equivalent, including, e.g., additional extracellular segments. The invention also includes a protein variation of the respective DCRS5 allele whose sequence is provided, e.g., a mutein or other construct. Typically, such variants will exhibit less than about 10% sequence differences with the target region, and thus will often have between 1- and 11-fold substitutions, e.g., 2-, 3-, 5, 7-fold, and others. It also encompasses allelic and other variants, e.g., natural polymorphisms, of the protein described. Typically, it will bind to its corresponding biological ligand, perhaps in a dimerized state with an alpha receptor subunit, with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10 nM, and more preferably at better than about 3 nM. The term shall also be used herein to refer to related naturally occurring forms, e.g., alleles, polymorphic variants, and metabolic variants of the mammalian protein. Preferred forms of the receptor complexes will bind the appropriate ligand with an affinity and selectivity appropriate for a ligand-receptor interaction.



This invention also encompasses combinations of proteins or peptides having substantial amino acid sequence identity with the amino acid sequence in Table 1. It will include sequence variants with relatively few substitutions, e.g., preferably fewer than about 3-5.

5       A substantial polypeptide "fragment", or "segment", is a stretch of amino acid residues of at least about 8 amino acids, generally at least 10 amino acids, more generally at least 12 amino acids, often at least 14 amino acids, more often at least 16 amino acids, typically at least 18 amino acids, more typically at least 20 amino acids, usually at least 22 amino acids, more usually at least 24 amino acids, preferably at least 26 amino acids, more preferably at least 28  
10   amino acids, and, in particularly preferred embodiments, at least about 30 or more amino acids. Sequences of segments of different proteins can be compared to one another over appropriate length stretches. In many situations, fragments may exhibit functional properties of the intact subunits, e.g., the extracellular domain of the transmembrane receptor may retain the ligand binding features, and may be used to prepare a soluble receptor-like complex.

15       Amino acid sequence homology, or sequence identity, is determined by optimizing residue matches. In some comparisons, gaps may be introduced, as required. See, e.g., Needleham, et al., (1970) J. Mol. Biol. 48:443-453; Sankoff, et al., (1983) chapter one in Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison, Addison-Wesley, Reading, MA; and software packages from IntelliGenetics, Mountain View,  
20   CA; and the University of Wisconsin Genetics Computer Group (GCG), Madison, WI; each of which is incorporated herein by reference. This changes when considering conservative substitutions as matches. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine.

25   Homologous amino acid sequences are intended to include natural allelic and interspecies variations in the cytokine sequence. Typical homologous proteins or peptides will have from 50 100% homology (if gaps can be introduced), to 60 100% homology (if conservative substitutions are included) with an amino acid sequence segment of Table 1. Homology measures will be at least about 70%, generally at least 76%, more generally at least 81%, often  
30   at least 85%, more often at least 88%, typically at least 90%, more typically at least 92%, usually at least 94%, more usually at least 95%, preferably at least 96%, and more preferably at least 97%, and in particularly preferred embodiments, at least 98% or more. The degree of homology will vary with the length of the compared segments. Homologous proteins or peptides, such as the allelic variants, will share most biological activities with the embodiments  
35   described in Table 1, particularly the intracellular portion.

As used herein, the term "biological activity" is used to describe, without limitation, effects on signaling, inflammatory responses, innate immunity, and/or morphogenic development by cytokine-like ligands. For example, these receptors should mediate phosphatase or phosphorylase activities, which activities are easily measured by standard procedures. See, e.g., Hardie, et al. (eds. 1995) *The Protein Kinase FactBook* vols. I and II, Academic Press, San Diego, CA; Hanks, et al. (1991) *Meth. Enzymol.* 200:38-62; Hunter, et al. (1992) *Cell* 70:375-388; Lewin (1990) *Cell* 61:743-752; Pines, et al. (1991) *Cold Spring Harbor Symp. Quant. Biol.* 56:449-463; and Parker, et al. (1993) *Nature* 363:736-738. The receptors, or portions thereof, may be useful as phosphate labeling enzymes to label general or specific substrates. The subunits may also be functional immunogens to elicit recognizing antibodies, or antigens capable of binding antibodies.

The terms ligand, agonist, antagonist, and analog of, e.g., a DCRS5 petition features of ligand-receptor interactions, e.g., where the receptor is a natural receptor or an antibody. The cellular responses likely are typically mediated through receptor tyrosine kinase pathways.

Also, a ligand is a molecule which serves either as a natural ligand to which said receptor, or an analog thereof, binds, or a molecule which is a functional analog of the natural ligand. The functional analog may be a ligand with structural modifications, or may be a wholly unrelated molecule which has a molecular shape which interacts with the appropriate ligand binding determinants. The ligands may serve as agonists or antagonists, see, e.g., Goodman, et al. (eds. 1990) *Goodman & Gilman's: The Pharmacological Bases of Therapeutics*, Pergamon Press, New York.

Rational drug design may also be based upon structural studies of the molecular shapes of a receptor or antibody and other effectors or ligands. See, e.g., Herz, et al. (1997) *J. Recept. Signal Transduct. Res.* 17:671-776; and Chaiken, et al. (1996) *Trends Biotechnol.* 14:369-375. Effectors may be other proteins which mediate other functions in response to ligand binding, or other proteins which normally interact with the receptor. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) *Protein Crystallography*, Academic Press, New York, which is hereby incorporated herein by reference.

## II. Activities

The cytokine receptor-like proteins will have a number of different biological activities, e.g., intracellular signaling, e.g., via STAT4, modulating cell proliferation, or in phosphate metabolism, being added to or removed from specific substrates, typically proteins. Such will

generally result in modulation of an inflammatory function, other innate immunity response, or a morphological effect. The subunit will probably have a specific low affinity binding to the ligand.

The DCRS5 has the characteristic motifs of a receptor signaling through the JAK pathway. See, e.g., Ihle, et al. (1997) *Stem Cells* 15(suppl. 1):105-111; Silvennoinen, et al. (1997) *APMIS* 105:497-509; Levy (1997) *Cytokine Growth Factor Review* 8:81-90; Winston and Hunter (1996) *Current Biol.* 6:668-671; Barrett (1996) *Baillieres Clin. Gastroenterol.* 10:1-15; and Briscoe, et al. (1996) *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 351:167-171. Of particular interest are the SH2 binding motifs described above.

The biological activities of the cytokine receptor subunits will be related to addition or removal of phosphate moieties to substrates, typically in a specific manner, but occasionally in a non specific manner. Substrates may be identified, or conditions for enzymatic activity may be assayed by standard methods, e.g., as described in Hardie, et al. (eds. 1995) *The Protein Kinase FactBook* vols. I and II, Academic Press, San Diego, CA; Hanks, et al. (1991) *Meth. Enzymol.* 200:38-62; Hunter, et al. (1992) *Cell* 70:375-388; Lewin (1990) *Cell* 61:743-752; Pines, et al. (1991) *Cold Spring Harbor Symp. Quant. Biol.* 56:449-463; and Parker, et al. (1993) *Nature* 363:736-738.

The receptor subunits may combine to form functional complexes, e.g., which may be useful for binding ligand or preparing antibodies. These will have substantial diagnostic uses, including detection or quantitation. The functional linkage of the receptor with the p40/IL-B30 ligand provides important insights into the clinical indications that the receptor will be useful for. Thus, antagonists and agonists will have predicted functional effects.

### III. Nucleic Acids

This invention contemplates use of isolated nucleic acid or fragments, e.g., which encode these or closely related proteins, or fragments thereof, e.g., to encode a corresponding polypeptide, preferably one which is biologically active. In addition, this invention covers isolated or recombinant DNAs which encode combinations of such proteins or polypeptides having characteristic sequences, e.g., of the DCRS5s alone or in combination with others such as an IL-12R $\beta$ 1 (see Showe, et al. (1996) *Ann. N.Y. Acad. Sci.* 795:413-425; Gately, et al. (1998) *Ann. Rev. Immunol.* 16:495-521; GenBank U03187, NM\_005535) subunit. Typically, the nucleic acid is capable of hybridizing, under appropriate conditions, with a nucleic acid sequence segment shown in Table 1, but preferably not with a corresponding segment of other receptors described in Table 3. Said biologically active protein or polypeptide can be a full length protein, or fragment, and will typically have a segment of amino acid sequence highly homologous, e.g., exhibiting significant stretches of identity, to one shown in Table 1. Further,

this invention covers the use of isolated or recombinant nucleic acid, or fragments thereof, which encode proteins having fragments which are equivalent to the DCRS5 proteins, e.g., intracellular portions. The isolated nucleic acids can have the respective regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition signals, and others from the natural gene. Combinations, as described, are also provided, e.g., combining the DCRS5 with the IL-12R $\beta$ 1, or their extracellular ligand binding portions as ligand antagonists. Diagnostic utilities are also clearly important, e.g., of polymorphic or other variants.

An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially pure, e.g., separated from other components which naturally accompany a native sequence, such as ribosomes, polymerases, and flanking genomic sequences from the originating species. The term embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates, which are thereby distinguishable from naturally occurring compositions, and chemically synthesized analogs or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated forms of the molecule, either completely or substantially pure.

An isolated nucleic acid will generally be a homogeneous composition of molecules, but will, in some embodiments, contain heterogeneity, preferably minor. This heterogeneity is typically found at the polymer ends or portions not critical to a desired biological function or activity.

A "recombinant" nucleic acid is typically defined either by its method of production or its structure. In reference to its method of production, e.g., a product made by a process, the process is use of recombinant nucleic acid techniques, e.g., involving human intervention in the nucleotide sequence. Typically this intervention involves in vitro manipulation, although under certain circumstances it may involve more classical animal breeding techniques. Alternatively, it can be a nucleic acid made by generating a sequence comprising fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally occurring mutants as found in their natural state. Thus, e.g., products made by transforming cells with an unnaturally occurring vector is encompassed, as are nucleic acids comprising sequence derived using any synthetic oligonucleotide process. Such a process is often done to replace, e.g., a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a restriction enzyme sequence recognition site, or for some structure-function analysis. Alternatively, the process is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms, e.g., encoding a fusion protein. Restriction enzyme recognition sites are often the target of such artificial manipulations, but other site specific targets, e.g.,

promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant, e.g., fusion, polypeptide. This will include a dimeric repeat or fusion of the DCRS5 with IL-12R $\beta$ 1 subunit. Specifically included are synthetic nucleic acids which, by genetic code redundancy,  
5 encode equivalent polypeptides to fragments of DCRS5 and fusions of sequences from various different related molecules, e.g., other cytokine receptor family members.

A "fragment" in a nucleic acid context is a contiguous segment of at least about 17 nucleotides, generally at least 21 nucleotides, more generally at least 25 nucleotides, ordinarily at least 30 nucleotides, more ordinarily at least 35 nucleotides, often at least 39 nucleotides,  
10 more often at least 45 nucleotides, typically at least 50 nucleotides, more typically at least 55 nucleotides, usually at least 60 nucleotides, more usually at least 66 nucleotides, preferably at least 72 nucleotides, more preferably at least 79 nucleotides, and in particularly preferred embodiments will be at least 85 or more nucleotides, including 90, 100, 120, 140, 160, 180, 200, etc. Typically, fragments of different genetic sequences can be compared to one another  
15 over appropriate length stretches, particularly defined segments such as the domains described below.

A nucleic acid which codes for the DCRS5 will be particularly useful to identify genes, mRNA, and cDNA species which code for itself or closely related proteins, as well as DNAs which code for polymorphic, allelic, or other genetic variants, e.g., from different individuals  
20 or related species. Preferred probes for such screens are those regions of the receptor which are conserved between different polymorphic variants or which contain nucleotides which lack specificity, and will preferably be full length or nearly so. In other situations, polymorphic variant specific sequences will be more useful. Combinations of polymorphic variants of DCRS5 with variants of IL-12R $\beta$ 1 may also be diagnosed.

25 This invention further covers recombinant nucleic acid molecules and fragments having a nucleic acid sequence identical to or highly homologous to the isolated DNA set forth herein. In particular, the sequences will often be operably linked to DNA segments which control transcription, translation, and DNA replication. These additional segments typically assist in expression of the desired nucleic acid segment.

30 Homologous, or highly identical, nucleic acid sequences, when compared to one another, e.g., DCRS5 sequences, exhibit significant similarity. The standards for homology in nucleic acids are either measures for homology generally used in the art by sequence comparison or based upon hybridization conditions. Comparative hybridization conditions are described in greater detail below.

35 Substantial identity in the nucleic acid sequence comparison context means either that the segments, or their complementary strands, when compared, are identical when optimally

aligned, with appropriate nucleotide insertions or deletions, in at least about 60% of the nucleotides, generally at least 66%, ordinarily at least 71%, often at least 76%, more often at least 80%, usually at least 84%, more usually at least 88%, typically at least 91%, more typically at least about 93%, preferably at least about 95%, more preferably at least about 96 to 5 98% or more, and in particular embodiments, as high at about 99% or more of the nucleotides, including, e.g., segments encoding structural domains or other segments described.

Alternatively, substantial identity will exist when the segments will hybridize under selective hybridization conditions, to a strand or its complement, typically using a sequence derived from Table 1. Typically, selective hybridization will occur when there is at least about 55% 10 homology over a stretch of at least about 14 nucleotides, more typically at least about 65%, preferably at least about 75%, and more preferably at least about 90%. See, Kanehisa (1984) Nucl. Acids Res. 12:203-213, which is incorporated herein by reference. The length of homology comparison, as described, may be over longer stretches, and in certain embodiments will be over a stretch of at least about 17 nucleotides, generally at least about 20 nucleotides, 15 ordinarily at least about 24 nucleotides, usually at least about 28 nucleotides, typically at least about 32 nucleotides, more typically at least about 40 nucleotides, preferably at least about 50 nucleotides, and more preferably at least about 75 to 100 or more nucleotides. This includes, e.g., 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, etc., and other lengths.

Stringent conditions, in referring to homology in the hybridization context, will be 20 stringent combined conditions of salt, temperature, organic solvents, and other parameters typically controlled in hybridization reactions. Stringent temperature conditions will usually include temperatures in excess of about 30° C, more usually in excess of about 37° C, typically in excess of about 45° C, more typically in excess of about 55° C, preferably in excess of about 65° C, and more preferably in excess of about 70° C. Stringent salt conditions will ordinarily 25 be less than about 500 mM, usually less than about 400 mM, more usually less than about 300 mM, typically less than about 200 mM, preferably less than about 100 mM, and more preferably less than about 80 mM, even down to less than about 50 or 20 mM. However, the combination of parameters is much more important than the measure of any single parameter. See, e.g., Wetmur and Davidson (1968) J. Mol. Biol. 31:349-370, which is hereby incorporated 30 herein by reference.

The isolated DNA can be readily modified by nucleotide substitutions, nucleotide deletions, nucleotide insertions, and inversions of nucleotide stretches. These modifications result in novel DNA sequences which encode this protein or its derivatives. These modified sequences can be used to produce mutant proteins (muteins) or to enhance the expression of 35 variant species. Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. Such mutant DCRS5 initiation of the DCRS5 as set

forth above, but having an amino acid sequence which differs from that of other cytokine receptor-like proteins as found in nature, whether by way of deletion, substitution, or insertion. In particular, "site specific mutant DCRS5" encompasses a protein having substantial sequence identity with a protein of Table 1, and typically shares most of the biological activities or effects of the forms disclosed herein. Various natural polymorphic variant sequences will also be identified.

Although site specific mutation sites are predetermined, mutants need not be site specific. Mammalian DCRS5 mutagenesis can be achieved by making amino acid insertions or deletions in the gene, coupled with expression. Substitutions, deletions, insertions, or many combinations may be generated to arrive at a final construct. Insertions include amino- or carboxy terminal fusions. Random mutagenesis can be conducted at a target codon and the expressed mammalian DCRS5 mutants can then be screened for the desired activity, providing some aspect of a structure-activity relationship. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis. See also Sambrook, et al. (1989) and Ausubel, et al. (1987 and periodic Supplements). Particularly useful constructs will be extracellular portions of the DCRS5 associated with IL-12R $\beta$ 1 segments.

The mutations in the DNA normally should not place coding sequences out of reading frames and preferably will not create complementary regions that could hybridize to produce secondary mRNA structure such as loops or hairpins.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Polymerase chain reaction (PCR) techniques can often be applied in mutagenesis. Alternatively, mutagenesis primers are commonly used methods for generating defined mutations at predetermined sites. See, e.g., Innis, et al. (eds. 1990) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA; and Dieffenbach and Dveksler (1995; eds.) PCR Primer: A Laboratory Manual Cold Spring Harbor Press, CSH, NY.

Certain embodiments of the invention are directed to combination compositions comprising the receptor sequences described. In other embodiments, functional portions of the sequences may be joined to encode fusion proteins. In other forms, variants of the described sequences may be substituted.

#### IV. Proteins, Peptides

As described above, the present invention encompasses primate DCRS5, e.g., whose sequences are disclosed in Table 1, and described above. Allelic and other variants are also contemplated, including, e.g., fusion proteins combining portions of such sequences with others, including, e.g., IL-12R $\beta$ 1, epitope tags, and functional domains.

The present invention also provides recombinant proteins, e.g., heterologous fusion proteins using segments from these primate or rodent proteins. A heterologous fusion protein is a fusion of proteins or segments which are naturally not normally fused in the same manner. Thus, the fusion product of a DCRS5 with another cytokine receptor is a continuous protein molecule having sequences fused in a typical peptide linkage, typically made as a single translation product and exhibiting properties, e.g., sequence or antigenicity, derived from each source peptide. A similar concept applies to heterologous nucleic acid sequences. Combinations of various designated proteins into complexes are also provided.

In addition, new constructs may be made from combining similar functional or structural domains from other related proteins, e.g., cytokine receptors or Toll-like receptors, including species variants. For example, ligand-binding or other segments may be "swapped" between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) Science 243:1330-1336; and O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992, each of which is incorporated herein by reference. Thus, new chimeric polypeptides exhibiting new combinations of specificities will result from the functional linkage of receptor-binding specificities. For example, the ligand binding domains from other related receptor molecules may be added or substituted for other domains of this or related proteins. The resulting protein will often have hybrid function and properties. For example, a fusion protein may include a targeting domain which may serve to provide sequestering of the fusion protein to a particular subcellular organelle.

Candidate fusion partners and sequences can be selected from various sequence data bases, e.g., GenBank, c/o IntelliGenetics, Mountain View, CA; and BCG, University of Wisconsin Biotechnology Computing Group, Madison, WI, which are each incorporated herein by reference. In particular, combinations of polypeptide sequences provided in Tables 1 and 3 are particularly preferred. Variant forms of the proteins may be substituted in the described combinations.

The present invention particularly provides muteins which bind cytokine-like ligands, and/or which are affected in signal transduction. Structural alignment of human DCRS5 with other members of the cytokine receptor family show conserved features/residues. See Table 3. Alignment of the human DCRS5 sequence with other members of the cytokine receptor family



indicates various structural and functionally shared features. See also, Bazan, et al. (1996) Nature 379:591; Lodi, et al. (1994) Science 263:1762-1766; Sayle and Milner-White (1995) TIBS 20:374-376; and Gronenberg, et al. (1991) Protein Engineering 4:263-269.

5 Substitutions with either mouse sequences or human sequences are particularly preferred. Conversely, conservative substitutions away from the ligand binding interaction regions will probably preserve most signaling activities; and conservative substitutions away from the intracellular domains will probably preserve most ligand binding properties.

"Derivatives" of the primate DCRS5 include amino acid sequence mutants, glycosylation variants, metabolic derivatives and covalent or aggregative conjugates with other  
10 chemical moieties. Covalent derivatives can be prepared by linkage of functionalities to groups which are found in the DCRS5 amino acid side chains or at the N termini, e.g., by means which are well known in the art. These derivatives can include, without limitation, aliphatic esters or amides of the carboxyl terminus, or of residues containing carboxyl side chains, O acyl derivatives of hydroxyl group containing residues, and N acyl derivatives of the  
15 amino terminal amino acid or amino group containing residues, e.g., lysine or arginine. Acyl groups are selected from the group of alkyl moieties, including C3 to C18 normal alkyl, thereby forming alkanoyl aroyl species.

In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further  
20 processing steps. Particularly preferred means for accomplishing this are by exposing the polypeptide to glycosylating enzymes derived from cells which normally provide such processing, e.g., mammalian glycosylation enzymes. Deglycosylation enzymes are also contemplated. Also embraced are versions of the same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues, e.g.,  
25 phosphotyrosine, phosphoserine, or phosphothreonine.

A major group of derivatives are covalent conjugates of the receptors or fragments thereof with other proteins of polypeptides. These derivatives can be synthesized in recombinant culture such as N terminal fusions or by the use of agents known in the art for their usefulness in cross linking proteins through reactive side groups. Preferred derivatization  
30 sites with cross linking agents are at free amino groups, carbohydrate moieties, and cysteine residues.

Fusion polypeptides between the receptors and other homologous or heterologous proteins are also provided. Homologous polypeptides may be fusions between different receptors, resulting in, for instance, a hybrid protein exhibiting binding specificity for multiple  
35 different cytokine ligands, or a receptor which may have broadened or weakened specificity of substrate effect. Likewise, heterologous fusions may be constructed which would exhibit a

combination of properties or activities of the derivative proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a receptor, e.g., a ligand-binding segment, so that the presence or location of a desired ligand may be easily determined. See, e.g., Dull, et al., U.S. Patent No. 4,859,609, which is hereby incorporated  
5 herein by reference. Other gene fusion partners include glutathione-S-transferase (GST), bacterial  $\beta$ -galactosidase, trpE, Protein A,  $\beta$ -lactamase, alpha amylase, alcohol dehydrogenase, and yeast alpha mating factor. See, e.g., Godowski, et al. (1988) Science 241:812-816. Labeled proteins will often be substituted in the described combinations of proteins. Associations of the DCRS5 with the IL-12R $\beta$ 1 are particularly significant, as described.

10 The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra, Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

15 Such polypeptides may also have amino acid residues which have been chemically modified by phosphorylation, sulfonation, biotinylation, or the addition or removal of other moieties, particularly those which have molecular shapes similar to phosphate groups. In some embodiments, the modifications will be useful labeling reagents, or serve as purification targets, e.g., affinity ligands.

20 Fusion proteins will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and expression are described generally, for example, in Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2d ed.), Vols. 1 3, Cold Spring Harbor Laboratory, and Ausubel, et al. (eds. 1987 and periodic supplements) Current Protocols in Molecular Biology, Greene/Wiley,  
25 New York, which are each incorporated herein by reference. Techniques for synthesis of polypeptides are described, for example, in Merrifield (1963) J. Amer. Chem. Soc. 85:2149-2156; Merrifield (1986) Science 232: 341-347; and Atherton, et al. (1989) Solid Phase Peptide Synthesis: A Practical Approach, IRL Press, Oxford; each of which is incorporated herein by reference. See also Dawson, et al. (1994) Science 266:776-779 for methods to make larger  
30 polypeptides.

This invention also contemplates the use of derivatives of a DCRS5 other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. These derivatives generally fall into three classes: (1) salts, (2) side chain and terminal residue covalent modifications, and (3) adsorption  
35 complexes, e.g., with cell membranes. Such covalent or aggregative derivatives are useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity

purification of a receptor or other binding molecule, e.g., an antibody. For example, a cytokine ligand can be immobilized by covalent bonding to a solid support such as cyanogen bromide activated Sepharose, by methods which are well known in the art, or adsorbed onto polyolefin surfaces, with or without glutaraldehyde cross linking, for use in the assay or purification of a  
5 cytokine receptor, antibodies, or other similar molecules. The ligand can also be labeled with a detectable group, e.g., radioiodinated by the chloramine T procedure, covalently bound to rare earth chelates, or conjugated to another fluorescent moiety for use in diagnostic assays.

A combination, e.g., including a DCRS5, of this invention can be used as an immunogen for the production of antisera or antibodies specific, e.g., capable of distinguishing  
10 between other cytokine receptor family members, for the combinations described. The complexes can be used to screen monoclonal antibodies or antigen-binding fragments prepared by immunization with various forms of impure preparations containing the protein. In particular, the term "antibodies" also encompasses antigen binding fragments of natural antibodies, e.g., Fab, Fab2, Fv, etc. The purified DCRS5 can also be used as a reagent to  
15 detect antibodies generated in response to the presence of elevated levels of expression, or immunological disorders which lead to antibody production to the endogenous receptor. Additionally, DCRS5 fragments may also serve as immunogens to produce the antibodies of the present invention, as described immediately below. For example, this invention contemplates antibodies having binding affinity to or being raised against the amino acid  
20 sequences shown in Table 1, fragments thereof, or various homologous peptides. In particular, this invention contemplates antibodies having binding affinity to, or having been raised against, specific fragments which are predicted to be, or actually are, exposed at the exterior protein surface of the native DCRS5. Complexes of combinations of proteins will also be useful, and antibody preparations thereto can be made.

25 In certain other embodiments, soluble constructs, e.g., of the extracellular ligand binding segments of the DCRS5 with the IL-12R $\beta$ 1 may be binding compositions for the ligand and may be useful as either ligand antagonists, or as antigens to block ligand mediated signaling. Such may be useful either diagnostically, e.g., for histology labeling for ligand, or therapeutically, e.g., as ligand antagonists.

30 The blocking of physiological response to the receptor ligands may result from the inhibition of binding of the ligand to the receptor, likely through competitive inhibition. Thus, in vitro assays of the present invention will often use antibodies or antigen binding segments of these antibodies, soluble receptor constructs, or fragments attached to solid phase substrates. These assays will also allow for the diagnostic determination of the effects of either ligand  
35 binding region mutations and modifications, or other mutations and modifications, e.g., which affect signaling or enzymatic function.

This invention also contemplates the use of competitive drug screening assays, e.g., where neutralizing antibodies to the receptor complexes or fragments compete with a test compound for binding to a ligand or other antibody. In this manner, the neutralizing antibodies or fragments can be used to detect the presence of a polypeptide which shares one or more binding sites to a receptor and can also be used to occupy binding sites on a receptor that might otherwise bind a ligand. Soluble receptor constructs combining the extracellular, or ligand binding, domains of the DCRS5 with the IL-12R $\beta$ 1, may be useful antagonists for competitive binding of p40/IL-B30 ligand.

#### 10 V. Making Nucleic Acids and Protein

DNA which encodes the protein or fragments thereof can be obtained by chemical synthesis, screening cDNA libraries, or by screening genomic libraries prepared from a wide variety of cell lines or tissue samples. Natural sequences can be isolated using standard methods and the sequences provided herein, e.g., in Table 1. Other species counterparts can be identified by hybridization techniques, or by various PCR techniques, combined with or by searching in sequence databases, e.g., GenBank.

This DNA can be expressed in a wide variety of host cells for the synthesis of a full length receptor or fragments which can in turn, for example, be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified ligand binding or kinase/phosphatase domains; and for structure/function studies. Variants or fragments can be expressed in host cells that are transformed or transfected with appropriate expression vectors. These molecules can be substantially free of protein or cellular contaminants, other than those derived from the recombinant host, and therefore are particularly useful in pharmaceutical compositions when combined with a pharmaceutically acceptable carrier and/or diluent. The protein, or portions thereof, may be expressed as fusions with other proteins. Combinations of the described proteins, or nucleic acids encoding them, are particularly interesting.

Expression vectors are typically self replicating DNA or RNA constructs containing the desired receptor gene, its fragments, or combination genes, usually operably linked to suitable genetic control elements that are recognized in a suitable host cell. These control elements are capable of effecting expression within a suitable host. Multiple genes may be coordinately expressed, and may be on a polycistronic message. The specific type of control elements necessary to effect expression will depend upon the eventual host cell used. Generally, the genetic control elements can include a prokaryotic promoter system or a eukaryotic promoter expression control system, and typically include a transcriptional promoter, an optional operator to control the onset of transcription, transcription enhancers to elevate the level of

mRNA expression, a sequence that encodes a suitable ribosome binding site, and sequences that terminate transcription and translation. Expression vectors also usually contain an origin of replication that allows the vector to replicate independently of the host cell.

5 The vectors of this invention include those which contain DNA which encodes a combination of proteins, as described, or a biologically active equivalent polypeptide. The DNA can be under the control of a viral promoter and can encode a selection marker. This invention further contemplates use of such expression vectors which are capable of expressing eukaryotic cDNAs coding for such proteins in a prokaryotic or eukaryotic host, where the vector is compatible with the host and where the eukaryotic cDNAs are inserted into the vector  
10 such that growth of the host containing the vector expresses the cDNAs in question. Usually, expression vectors are designed for stable replication in their host cells or for amplification to greatly increase the total number of copies of the desirable gene(s) per cell. It is not always necessary to require that an expression vector replicate in a host cell, e.g., it is possible to effect transient expression of the protein or its fragments in various hosts using vectors that do not  
15 contain a replication origin that is recognized by the host cell. It is also possible to use vectors that cause integration of the protein encoding portions into the host DNA by recombination.

Vectors, as used herein, comprise plasmids, viruses, bacteriophage, integratable DNA fragments, and other vehicles which enable the integration of DNA fragments into the genome of the host. Expression vectors are specialized vectors which contain genetic control elements  
20 that effect expression of operably linked genes. Plasmids are the most commonly used form of vector but all other forms of vectors which serve an equivalent function and which are, or become, known in the art are suitable for use herein. See, e.g., Pouwels, et al. (1985 and Supplements) Cloning Vectors: A Laboratory Manual, Elsevier, N.Y., and Rodriguez, et al. (eds. 1988) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworth,  
25 Boston, which are incorporated herein by reference.

Transformed cells are cells, preferably mammalian, that have been transformed or transfected with vectors constructed using recombinant DNA techniques. Transformed host cells usually express the desired proteins, but for purposes of cloning, amplifying, and manipulating its DNA, do not need to express the subject proteins. This invention further  
30 contemplates culturing transformed cells in a nutrient medium, thus permitting the proteins to accumulate. The proteins can be recovered, either from the culture or, in certain instances, from the culture medium.

For purposes of this invention, nucleic sequences are operably linked when they are functionally related to each other. For example, DNA for a presequence or secretory leader is  
35 operably linked to a polypeptide if it is expressed as a preprotein or participates in directing the polypeptide to the cell membrane or in secretion of the polypeptide. A promoter is operably

linked to a coding sequence if it controls the transcription of the polypeptide; a ribosome binding site is operably linked to a coding sequence if it is positioned to permit translation. Usually, operably linked means contiguous and in reading frame, however, certain genetic elements such as repressor genes are not contiguously linked but still bind to operator sequences that in turn control expression.

Suitable host cells include prokaryotes, lower eukaryotes, and higher eukaryotes. Prokaryotes include both gram negative and gram positive organisms, e.g., *E. coli* and *B. subtilis*. Lower eukaryotes include yeasts, e.g., *S. cerevisiae* and *Pichia*, and species of the genus *Dictyostelium*. Higher eukaryotes include established tissue culture cell lines from animal cells, both of non mammalian origin, e.g., insect cells, and birds, and of mammalian origin, e.g., human, primates, and rodents.

Prokaryotic host vector systems include a wide variety of vectors for many different species. As used herein, *E. coli* and its vectors will be used generically to include equivalent vectors used in other prokaryotes. A representative vector for amplifying DNA is pBR322 or many of its derivatives. Vectors that can be used to express the receptor or its fragments include, but are not limited to, such vectors as those containing the lac promoter (pUC series); trp promoter (pBR322 trp); Ipp promoter (the pIN series); lambda pP or pR promoters (pOTS); or hybrid promoters such as ptac (pDR540). See Brosius, et al. (1988) "Expression Vectors Employing Lambda, and Ipp derived Promoters", in *Vectors: A Survey of Molecular Cloning Vectors and Their Uses*, (eds. Rodriguez and Denhardt), Butterworth, Boston, Chapter 10, pp. 205-236, which is incorporated herein by reference.

Lower eukaryotes, e.g., yeasts and *Dictyostelium*, may be transformed with DCRS5 sequence containing vectors. For purposes of this invention, the most common lower eukaryotic host is the baker's yeast, *Saccharomyces cerevisiae*. It will be used to generically represent lower eukaryotes although a number of other strains and species are also available. Yeast vectors typically consist of a replication origin (unless of the integrating type), a selection gene, a promoter, DNA encoding the receptor or its fragments, and sequences for translation termination, polyadenylation, and transcription termination. Suitable expression vectors for yeast include such constitutive promoters as 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionein promoter. Suitable vectors include derivatives of the following types: self-replicating low copy number (such as the YRp series), self-replicating high copy number (such as the YEpl series); integrating types (such as the YIp series), or mini-chromosomes (such as the YCp series).

Higher eukaryotic tissue culture cells are normally the preferred host cells for expression of the functionally active interleukin or receptor proteins. In principle, many higher

eukaryotic tissue culture cell lines are workable, e.g., insect baculovirus expression systems, whether from an invertebrate or vertebrate source. However, mammalian cells are preferred. Transformation or transfection and propagation of such cells has become a routine procedure. Examples of useful cell lines include HeLa cells, Chinese hamster ovary (CHO) cell lines, baby rat kidney (BRK) cell lines, insect cell lines, bird cell lines, and monkey (COS) cell lines. Expression vectors for such cell lines usually include an origin of replication, a promoter, a translation initiation site, RNA splice sites (if genomic DNA is used), a polyadenylation site, and a transcription termination site. These vectors also usually contain a selection gene or amplification gene. Suitable expression vectors may be plasmids, viruses, or retroviruses carrying promoters derived, e.g., from such sources as from adenovirus, SV40, parvoviruses, vaccinia virus, or cytomegalovirus. Representative examples of suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) *Mol. Cell Biol.* 5:1136-1142; pMC1neo PolyA, see Thomas, et al. (1987) *Cell* 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610.

For secreted proteins and some membrane proteins, an open reading frame usually encodes a polypeptide that consists of a mature or secreted product covalently linked at its N-terminus to a signal peptide. The signal peptide is cleaved prior to secretion of the mature, or active, polypeptide. The cleavage site can be predicted with a high degree of accuracy from empirical rules, e.g., von-Heijne (1986) *Nucleic Acids Research* 14:4683-4690 and Nielsen, et al. (1997) *Protein Eng.* 10:1-12, and the precise amino acid composition of the signal peptide often does not appear to be critical to its function, e.g., Randall, et al. (1989) *Science* 243:1156-1159; Kaiser et al. (1987) *Science* 235:312-317. The mature proteins of the invention can be readily determined using standard methods.

It will often be desired to express these polypeptides in a system which provides a specific or defined glycosylation pattern. In this case, the usual pattern will be that provided naturally by the expression system. However, the pattern will be modifiable by exposing the polypeptide, e.g., an unglycosylated form, to appropriate glycosylating proteins introduced into a heterologous expression system. For example, the receptor gene may be co-transformed with one or more genes encoding mammalian or other glycosylating enzymes. Using this approach, certain mammalian glycosylation patterns will be achievable in prokaryote or other cells. Expression in prokaryote cells will typically lead to unglycosylated forms of protein.

The source of DCRS5 can be a eukaryotic or prokaryotic host expressing recombinant DCRS5, such as is described above. The source can also be a cell line, but other mammalian cell lines are also contemplated by this invention, with the preferred cell line being from the human species.

Now that the sequences are known, the primate DCRS5, fragments, or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These include processes such as are described in Stewart and Young (1984) Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, IL; Bodanszky and Bodanszky (1984) The Practice of Peptide  
5 Synthesis, Springer Verlag, New York; and Bodanszky (1984) The Principles of Peptide Synthesis, Springer Verlag, New York; all of each which are incorporated herein by reference. For example, an azide process, an acid chloride process, an acid anhydride process, a mixed anhydride process, an active ester process (for example, p nitrophenyl ester, N hydroxysuccinimide ester, or cyanomethyl ester), a carbodiimidazole process, an oxidative  
10 reductive process, or a dicyclohexylcarbodiimide (DCCD) additive process can be used. Solid phase and solution phase syntheses are both applicable to the foregoing processes. Similar techniques can be used with partial DCRS5 sequences.

The DCRS5 proteins, fragments, or derivatives are suitably prepared in accordance with the above processes as typically employed in peptide synthesis, generally either by a so  
15 called stepwise process which comprises condensing an amino acid to the terminal amino acid, one by one in sequence, or by coupling peptide fragments to the terminal amino acid. Amino groups that are not being used in the coupling reaction typically must be protected to prevent coupling at an incorrect location.

If a solid phase synthesis is adopted, the C terminal amino acid is bound to an insoluble  
20 carrier or support through its carboxyl group. The insoluble carrier is not particularly limited as long as it has a binding capability to a reactive carboxyl group. Examples of such insoluble carriers include halomethyl resins, such as chloromethyl resin or bromomethyl resin, hydroxymethyl resins, phenol resins, tert alkylloxycarbonylhydrazidated resins, and the like.

An amino group protected amino acid is bound in sequence through condensation of its  
25 activated carboxyl group and the reactive amino group of the previously formed peptide or chain, to synthesize the peptide step by step. After synthesizing the complete sequence, the peptide is split off from the insoluble carrier to produce the peptide. This solid phase approach is generally described by Merrifield, et al. (1963) in J. Am. Chem. Soc. 85:2149 2156, which is incorporated herein by reference.

30 The prepared protein and fragments thereof can be isolated and purified from the reaction mixture by means of peptide separation, e.g., by extraction, precipitation, electrophoresis, various forms of chromatography, immunoaffinity, and the like. The receptors of this invention can be obtained in varying degrees of purity depending upon desired uses. Purification can be accomplished by use of the protein purification techniques disclosed herein,  
35 see below, or by the use of the antibodies herein described in methods of immunoabsorbant affinity chromatography. This immunoabsorbant affinity chromatography is carried out by



first linking the antibodies to a solid support and then contacting the linked antibodies with solubilized lysates of appropriate cells, lysates of other cells expressing the receptor, or lysates or supernatants of cells producing the protein as a result of DNA techniques, see below.

5 Generally, the purified protein will be at least about 40% pure, ordinarily at least about 50% pure, usually at least about 60% pure, typically at least about 70% pure, more typically at least about 80% pure, preferable at least about 90% pure and more preferably at least about 95% pure, and in particular embodiments, 97%-99% or more. Purity will usually be on a weight basis, but can also be on a molar basis. Different assays will be applied as appropriate. Individual proteins may be purified and thereafter combined.

10

## VI. Antibodies

Antibodies can be raised to the various mammalian, e.g., primate DCRS5 proteins and fragments thereof, both in naturally occurring native forms and in their recombinant forms, the difference being that antibodies to the active receptor are more likely to recognize epitopes  
15 which are only present in the native conformations. Antibodies recognizing epitopes presented by the combination, e.g., functionally, of the DCRS5 with the IL-12R $\beta$ 1 are also contemplated. Denatured antigen detection can also be useful in, e.g., Western analysis. Anti-idiotypic antibodies are also contemplated, which would be useful as agonists or antagonists of a natural receptor or an antibody.

20 Antibodies, including binding fragments and single chain versions, against predetermined fragments of the protein can be raised by immunization of animals with conjugates of the fragments with immunogenic proteins. Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective protein, or screened for agonistic or antagonistic activity. These  
25 monoclonal antibodies will usually bind with at least a KD of about 1 mM, more usually at least about 300  $\mu$ M, typically at least about 100  $\mu$ M, more typically at least about 30  $\mu$ M, preferably at least about 10  $\mu$ M, and more preferably at least about 3  $\mu$ M or better.

The antibodies, including antigen binding fragments, of this invention can have significant diagnostic or therapeutic value. They can be potent antagonists that bind to the  
30 receptor and inhibit binding to ligand or inhibit the ability of the receptor to elicit a biological response, e.g., act on its substrate. They also can be useful as non neutralizing antibodies and can be coupled to toxins or radionuclides to bind producing cells, or cells localized to the source of the interleukin. Further, these antibodies can be conjugated to drugs or other therapeutic agents, either directly or indirectly by means of a linker.

35 The antibodies of this invention can also be useful in diagnostic applications. As capture or non neutralizing antibodies, they might bind to the receptor without inhibiting ligand

or substrate binding. As neutralizing antibodies, they can be useful in competitive binding assays. They will also be useful in detecting or quantifying ligand. They may be used as reagents for Western blot analysis, or for immunoprecipitation or immunopurification of the respective protein. Likewise, nucleic acids and proteins may be immobilized to solid  
5 substrates for affinity purification or detection methods. The substrates may be, e.g., solid resin beads or sheets of plastic.

Protein fragments may be joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as immunogens. Mammalian cytokine receptors and fragments may be fused or covalently linked to a variety of immunogens, such as keyhole  
10 limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See Microbiology, Hoeber Medical Division, Harper and Row, 1969; Landsteiner (1962) Specificity of Serological Reactions, Dover Publications, New York; and Williams, et al. (1967) Methods in Immunology and Immunochemistry, Vol. 1, Academic Press, New York; each of which are incorporated herein by reference, for descriptions of methods of preparing polyclonal antisera.  
15 A typical method involves hyperimmunization of an animal with an antigen. The blood of the animal is then collected shortly after the repeated immunizations and the gamma globulin is isolated.

In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as mice, rodents, primates, humans, etc. Description of techniques for  
20 preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds.) Basic and Clinical Immunology (4th ed.), Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH Press; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed.) Academic Press, New York; and particularly in Kohler and Milstein (1975) in Nature 256: 495 497, which discusses one  
25 method of generating monoclonal antibodies. Each of these references is incorporated herein by reference. Summarized briefly, this method involves injecting an animal with an immunogen. The animal is then sacrificed and cells taken from its spleen, which are then fused with myeloma cells. The result is a hybrid cell or "hybridoma" that is capable of reproducing in vitro. The population of hybridomas is then screened to isolate individual  
30 clones, each of which secrete a single antibody species to the immunogen. In this manner, the individual antibody species obtained are the products of immortalized and cloned single B cells from the immune animal generated in response to a specific site recognized on the immunogenic substance.

Other suitable techniques involve in vitro exposure of lymphocytes to the antigenic  
35 polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors. See, Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin

Repertoire in Phage Lambda," Science 246:1275-1281; and Ward, et al. (1989) Nature 341:544-546, each of which is hereby incorporated herein by reference. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents, teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant or chimeric immunoglobulins may be produced, see Cabilly, U.S. Patent No. 4,816,567; or made in transgenic mice, see Mendez, et al. (1997) Nature Genetics 15:146-156. These references are incorporated herein by reference.

The antibodies of this invention can also be used for affinity chromatography in isolating the DCRS5 proteins or peptides. Columns can be prepared where the antibodies are linked to a solid support, e.g., particles, such as agarose, Sephadex, or the like, where a cell lysate may be passed through the column, the column washed, followed by increasing concentrations of a mild denaturant, whereby the purified protein will be released. Alternatively, the protein may be used to purify antibody. Appropriate cross absorptions or depletions may be applied.

The antibodies may also be used to screen expression libraries for particular expression products. Usually the antibodies used in such a procedure will be labeled with a moiety allowing easy detection of presence of antigen by antibody binding.

Antibodies raised against a cytokine receptor will also be used to raise anti-idiotypic antibodies. These will be useful in detecting or diagnosing various immunological conditions related to expression of the protein or cells which express the protein. They also will be useful as agonists or antagonists of the ligand, which may be competitive receptor inhibitors or substitutes for naturally occurring ligands. Certain antibodies to receptor subunits or combinations may serve as activating antibodies, which may effect signaling thereby serving, e.g., as ligand agonists.

A cytokine receptor protein that specifically binds to or that is specifically immunoreactive with an antibody generated against a defined immunogen, such as an immunogen consisting of the amino acid sequence of SEQ ID NO: 2, is typically determined in an immunoassay. The immunoassay typically uses a polyclonal antiserum which was raised, e.g., to a protein of SEQ ID NO: 2. This antiserum is selected to have low crossreactivity against other cytokine receptor family members, e.g., IL-12R $\beta$ 2 receptor subunit or IL-6

receptor subunit gp130, preferably from the same species, and any such crossreactivity is removed by immunoabsorption prior to use in the immunoassay.

5 In order to produce antisera for use in an immunoassay, the protein, e.g., of SEQ ID NO: 2, is isolated as described herein. For example, recombinant protein may be produced in a mammalian cell line. An appropriate host, e.g., an inbred strain of mice such as Balb/c, is immunized with the selected protein, typically using a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see Harlow and Lane, *supra*). Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used as an immunogen. Polyclonal sera are collected and titrated  
10 against the immunogen protein in an immunoassay, e.g., a solid phase immunoassay with the immunogen immobilized on a solid support. Polyclonal antisera with a titer of  $10^4$  or greater are selected and tested for their cross reactivity against other cytokine receptor family members, e.g., gp130 or IL-12R $\beta$ 1 using a competitive binding immunoassay such as the one described in Harlow and Lane, *supra*, at pages 570-573. Preferably at least two cytokine  
15 receptor family members are used in this determination. These cytokine receptor family members can be produced as recombinant proteins and isolated using standard molecular biology and protein chemistry techniques as described herein.

Immunoassays in the competitive binding format can be used for the crossreactivity determinations. For example, the protein of SEQ ID NO: 2 can be immobilized to a solid  
20 support. Proteins added to the assay compete with the binding of the antisera to the immobilized antigen. The ability of the above proteins to compete with the binding of the antisera to the immobilized protein is compared to the proteins, e.g., of gp130 or IL-12R $\beta$ 2. The percent crossreactivity for the above proteins is calculated, using standard calculations. Those antisera with less than 10% crossreactivity with each of the proteins listed above are  
25 selected and pooled. The cross-reacting antibodies are then removed from the pooled antisera by immunoabsorption with the above-listed proteins.

The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein to the immunogen protein (e.g., the DCRS5 like protein of SEQ ID NO: 2). In order to make this comparison, the two proteins  
30 are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required is less than twice the amount of the protein of the selected protein or proteins that is required, then the second protein is said to specifically bind to an antibody generated to the immunogen.

35 It is understood that these cytokine receptor proteins are members of a family of homologous proteins that comprise many identified genes. For a particular gene product, such

as the DCRS5, the term refers not only to the amino acid sequences disclosed herein, but also to other proteins that are allelic, non-allelic, or species variants. It is also understood that the terms include nonnatural mutations introduced by deliberate mutation using conventional recombinant technology such as single site mutation, or by excising short sections of DNA encoding the respective proteins, or by substituting new amino acids, or adding new amino acids. Such minor alterations typically will substantially maintain the immunoidentity of the original molecule and/or its biological activity. Thus, these alterations include proteins that are specifically immunoreactive with a designated naturally occurring DCRS5 protein. The biological properties of the altered proteins can be determined by expressing the protein in an appropriate cell line and measuring the appropriate effect, e.g., upon transfected lymphocytes. Particular protein modifications considered minor would include conservative substitution of amino acids with similar chemical properties, as described above for the cytokine receptor family as a whole. By aligning a protein optimally with the protein of the cytokine receptors and by using the conventional immunoassays described herein to determine immunoidentity, one can determine the protein compositions of the invention.

Moreover, antibodies against the receptor subunits may serve to sterically block ligand binding to the functional receptor. Such antibodies may be raised to either subunit alone, or to the combination of DCRS5 with IL-12R $\beta$ 1. Antibody antagonists would result.

## VII. Kits, diagnosis, and quantitation

Both naturally occurring and recombinant forms of the cytokine receptor like molecules of this invention are particularly useful in kits and assay methods. For example, these methods would also be applied to screening for binding activity, e.g., ligands for these proteins. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds per year. See, e.g., a BIOMEK automated workstation, Beckman Instruments, Palo Alto, California, and Fodor, et al. (1991) Science 251:767-773, which is incorporated herein by reference. The latter describes means for testing binding by a plurality of defined polymers synthesized on a solid substrate. The development of suitable assays to screen for a ligand or agonist/antagonist homologous proteins can be greatly facilitated by the availability of large amounts of purified, soluble cytokine receptors in an active state such as is provided by this invention.

Purified DCRS5 can be coated directly onto plates for use in the aforementioned ligand screening techniques. However, non neutralizing antibodies to these proteins can be used as capture antibodies to immobilize the respective receptor on the solid phase, useful, e.g., in diagnostic uses.

This invention also contemplates use of DCRS5, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for detecting the presence of the protein or its ligand. Alternatively, or additionally, antibodies against the molecules may be incorporated into the kits and methods. Typically the kit will have a compartment containing  
5 either a DCRS5 peptide or gene segment or a reagent which recognizes one or the other. Typically, recognition reagents, in the case of peptide, would be a receptor or antibody, or in the case of a gene segment, would usually be a hybridization probe. Other kit components may include other proteins or reagents related to the p40, IL-B30, or IL-12R $\beta$ 1 polypeptides of the ligand/receptor pairing.

10 A preferred kit for determining the concentration of DCRS5 in a sample would typically comprise a labeled compound, e.g., ligand or antibody, having known binding affinity for DCRS5, a source of DCRS5 (naturally occurring or recombinant) as a positive control, and a means for separating the bound from free labeled compound, for example a solid phase for immobilizing the DCRS5 in the test sample. Compartments containing reagents, and  
15 instructions, will normally be provided. Appropriate nucleic acid or protein containing kits are also provided.

Antibodies, including antigen binding fragments, specific for mammalian DCRS5 or a peptide fragment, or receptor fragments are useful in diagnostic applications to detect the presence of elevated levels of ligand and/or its fragments. Diagnostic assays may be  
20 homogeneous (without a separation step between free reagent and antibody-antigen complex) or heterogeneous (with a separation step). Various commercial assays exist, such as radioimmunoassay (RIA), enzyme linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme multiplied immunoassay technique (EMIT), substrate labeled fluorescent immunoassay (SLFIA) and the like. For example, unlabeled antibodies can be  
25 employed by using a second antibody which is labeled and which recognizes the antibody to a cytokine receptor or to a particular fragment thereof. These assays have also been extensively discussed in the literature. See, e.g., Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH., and Coligan (ed. 1991 and periodic supplements) Current Protocols In Immunology Greene/Wiley, New York.

30 Anti-idiotypic antibodies may have similar use to serve as agonists or antagonists of cytokine receptors. These should be useful as therapeutic reagents under appropriate circumstances.

Frequently, the reagents for diagnostic assays are supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the  
35 protocol, and the label, either labeled or unlabeled antibody, or labeled ligand is provided. This is usually in conjunction with other additives, such as buffers, stabilizers, materials

necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also contain instructions for proper use and disposal of the contents after use. Typically the kit has compartments for each useful reagent, and will contain instructions for proper use and disposal of reagents. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium having appropriate concentrations for performing the assay.

The aforementioned constituents of the diagnostic assays may be used without modification or may be modified in a variety of ways. For example, labeling may be achieved by covalently or non covalently joining a moiety which directly or indirectly provides a detectable signal. In many of these assays, a test compound, cytokine receptor, or antibodies thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as  $^{125}\text{I}$ , enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Both of the patents are incorporated herein by reference. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

There are also numerous methods of separating the bound from the free ligand, or alternatively the bound from the free test compound. The cytokine receptor can be immobilized on various matrixes followed by washing. Suitable matrices include plastic such as an ELISA plate, filters, and beads. Methods of immobilizing the receptor to a matrix include, without limitation, direct adhesion to plastic, use of a capture antibody, chemical coupling, and biotin avidin. The last step in this approach involves the precipitation of antibody/antigen complex by any of several methods including those utilizing, e.g., an organic solvent such as polyethylene glycol or a salt such as ammonium sulfate. Other suitable separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) Clin. Chem. 30(9):1457-1461, and the double antibody magnetic particle separation as described in U.S. Pat. No. 4,659,678, each of which is incorporated herein by reference.

The methods for linking protein or fragments to various labels have been extensively reported in the literature and do not require detailed discussion here. Many of the techniques involve the use of activated carboxyl groups either through the use of carbodiimide or active esters to form peptide bonds, the formation of thioethers by reaction of a mercapto group with an activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

Another diagnostic aspect of this invention involves use of oligonucleotide or polynucleotide sequences taken from the sequence of an cytokine receptor. These sequences can be used as probes for detecting levels of the respective cytokine receptor in patients suspected of having an immunological disorder. The preparation of both RNA and DNA  
5 nucleotide sequences, the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. Normally an oligonucleotide probe should have at least about 14 nucleotides, usually at least about 18 nucleotides, and the polynucleotide probes may be up to several kilobases. Various labels may be employed, most commonly radionuclides, particularly  $^{32}\text{P}$ . However, other techniques may also be employed,  
10 such as using biotin modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides, fluorescers, enzymes, or the like. Alternatively, antibodies may be employed which can recognize specific duplexes, including DNA duplexes, RNA duplexes, DNA RNA hybrid duplexes, or DNA protein duplexes. The antibodies in turn  
15 may be labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected. The use of probes to the novel anti sense RNA may be carried out in conventional techniques such as nucleic acid hybridization, plus and minus screening, recombinational probing, hybrid released translation (HRT), and hybrid arrested translation (HART). This also  
20 includes amplification techniques such as polymerase chain reaction (PCR).

Diagnostic kits which also test for the qualitative or quantitative presence of other markers are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers. Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) Progress in Growth Factor Res. 1:89-97. Detection of polymorphic  
25 variations, which may reflect functional receptor signaling differences, may be useful in determining therapeutic strategy. Variations which reflect greater or lesser response to ligand may allow subsetting of responsive/non-responsive patient pools.

#### VIII. Therapeutic Utility

30 This invention provides reagents with significant therapeutic value. See, e.g., Levitzki (1996) Curr. Opin. Cell Biol. 8:239-244. The cytokine receptors (naturally occurring or recombinant), fragments thereof, mutein receptors, and antibodies, along with compounds identified as having binding affinity to the receptors or antibodies, should be useful in the treatment of conditions exhibiting abnormal expression of the receptors or their ligands. Such  
35 abnormality will typically be manifested by immunological disorders. See WO 01/18051, which is incorporated herein by reference. Additionally, this invention should provide



therapeutic value in various diseases or disorders associated with abnormal expression or abnormal triggering of response to the ligand. For example, the p40/IL B30 ligand has been suggested to be involved in development of cell mediated immunity, e.g., anti-tumor activity, mounting of humoral and cellular immunity, and antiviral effects. In particular, the ligand  
5 appears to activate NK and T cells. Therapy may be combined with IL-18, IL-12, TNF, IFN $\gamma$ , radiation/chemo therapy, adjuvants, or antitumor, antiviral, or antifungal compounds.

Conversely, antagonists, which may be combined with antagonists of TNF, IFN $\gamma$ , IL-18, or IL-12, or with IL-10 or steroids, may be indicated in chronic Th1 mediated diseases, autoimmunity, or transplant and/or rejection situations, multiple sclerosis, psoriasis, chronic  
10 inflammatory conditions, rheumatoid arthritis, osteoarthritis, or inflammatory bowel diseases. Antagonists may take the form of antibodies against the receptor subunits, soluble receptor constructs, or antisense nucleic acids to one or more of the the receptor subunits. The matching of the p40/IL-B30 ligand with receptor subunits DCRS5 and IL-12R $\beta$ 1 provides insight into indications for use of the agonists and antagonists.

15 Therapeutically, based on the p40/IL-B30 activities described, antagonists of the cytokine may be effected, e.g., by soluble DCRS5, with or without soluble IL-12R $\beta$ 1, or antibodies to either receptor subunit. Antagonists may be useful as inhibitors of undesirable immune or inflammatory responses, to target memory T cells, or in combination with IL-12/IL-12R antagonists, or other anti-inflammatories or immunosuppressants. Clinical indications  
20 may be chronic inflammation or transplant situations. Various polymorphisms may enhance or decrease receptor function, and if dominant, might be useful as therapeutics. Identification of such variants may allow subsetting of responsive or nonresponsive patient pools. The reagents may be useful as detecting or labeling reagents or ablative reagents for memory T cells and/or NK cells.

25 Gene therapy may render desired cell populations response to p40/IL-B30 ligand, e.g., as adjuvants for tumor immunotherapy, to facilitate activation of tumor infiltrating lymphocytes, T cells, or NK cells. Antisense strategies may be applied, e.g., to prevent receptor responsiveness.

Various abnormal conditions are known in cell types shown to produce both IL-12 p40  
30 and/or IL-B30 mRNA by Northern blot analysis. See Berkow (ed.) The Merck Manual of Diagnosis and Therapy, Merck & Co., Rahway, N.J.; Thorn, et al. Harrison's Principles of Internal Medicine, McGraw-Hill, N.Y.; and Weatherall, et al. (eds.) Oxford Textbook of Medicine, Oxford University Press, Oxford. Many other medical conditions and diseases will be responsive to treatment by an agonist or antagonist provided herein. See, e.g., Stites and  
35 Terr (eds.; 1991) Basic and Clinical Immunology Appleton and Lange, Norwalk, Connecticut; and Samter, et al. (eds.) Immunological Diseases Little, Brown and Co. Other likely

indications for treatment include bone remodeling, sexual dysfunction, prevention of neurodegenerative diseases, dementia, stress, and others. These problems should be susceptible to prevention or treatment using compositions provided herein.

5 Recombinant cytokine receptors, muteins, agonist or antagonist antibodies thereto, or antibodies can be purified and then administered to a patient. These reagents can be combined for therapeutic use with additional active ingredients, e.g., in conventional pharmaceutically acceptable carriers or diluents, along with physiologically innocuous stabilizers and excipients. These combinations can be sterile, e.g., filtered, and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also  
10 contemplates use of antibodies or binding fragments thereof which are not complement binding.

Ligand screening using cytokine receptor or fragments thereof can be performed to identify molecules having binding affinity to the receptors. Subsequent biological assays can then be utilized to determine if a putative ligand can provide competitive binding, which can  
15 block intrinsic stimulating activity. Receptor fragments can be used as a blocker or antagonist in that it blocks the activity of ligand. Likewise, a compound having intrinsic stimulating activity can activate the receptor and is thus an agonist in that it simulates the activity of ligand, e.g., inducing signaling. This invention further contemplates the therapeutic use of antibodies to cytokine receptors as antagonists.

20 The quantities of reagents necessary for effective therapy will depend upon many different factors, including means of administration, target site, reagent physiological life, pharmacological life, physiological state of the patient, and other medicants administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts useful for in situ administration of  
25 these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication of human dosage. Various considerations are described, e.g., in Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn.; each of which is hereby incorporated herein by  
30 reference. Methods for administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others. Pharmaceutically acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, New Jersey. Because of the likely high affinity binding, or turnover numbers, between a putative ligand and its receptors, low  
35 dosages of these reagents would be initially expected to be effective. And the signaling pathway suggests extremely low amounts of ligand may have effect. Thus, dosage ranges

would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10  $\mu$ M concentrations, usually less than about 100 nM, preferably less than about 10 pM (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or slow release apparatus will often be utilized for continuous administration.

Cytokine receptors, fragments thereof, and antibodies or its fragments, antagonists, and agonists, may be administered directly to the host to be treated or, depending on the size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in many conventional dosage formulations. While it is possible for the active ingredient to be administered alone, it is preferable to present it as a pharmaceutical formulation. Formulations comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier must be both pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient.

Formulations include those suitable for oral, rectal, nasal, or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn.; Avis, et al. (eds. 1993) Pharmaceutical Dosage Forms: Parenteral Medications Dekker, NY; Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Tablets Dekker, NY; and Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Disperse Systems Dekker, NY. The therapy of this invention may be combined with or used in association with other therapeutic agents, particularly agonists or antagonists of other cytokine receptor family members.

## IX. Screening

Drug screening using DCRS5 or fragments thereof can be performed to identify compounds having binding affinity to the receptor subunit, including isolation of associated components. Subsequent biological assays can then be utilized to determine if the compound has intrinsic stimulating activity and is therefore a blocker or antagonist in that it blocks the activity of the ligand.

Moreover, matching of the p40/IL-B30 ligand with the functional receptor of DCRS3 with IL-12R $\beta$ 1, allows screening for antagonists and agonists with a positive signaling control. Small molecule or antibody screening can be done.

One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant DNA molecules expressing the DCRS5 in combination with another cytokine receptor subunit, e.g., the IL-12R $\beta$ 1. The signaling is believed to use STAT4. Cells may be isolated which express a receptor in isolation from other functional receptors. Such cells, either in viable or fixed form, can be used for standard antibody/antigen or ligand/receptor binding assays. See also, Parce, et al. (1989) *Science* 246:243-247; and Owicki, et al. (1990) *Proc. Nat'l Acad. Sci. USA* 87:4007-4011, which describe sensitive methods to detect cellular responses. Competitive assays are particularly useful, where the cells are contacted and incubated with a labeled receptor or antibody having known binding affinity to the ligand, such as <sup>125</sup>I-antibody, and a test sample whose binding affinity to the binding composition is being measured. The bound and free labeled binding compositions are then separated to assess the degree of ligand binding. The amount of test compound bound is inversely proportional to the amount of labeled receptor binding to the known source. Many techniques can be used to separate bound from free ligand to assess the degree of ligand binding. This separation step could typically involve a procedure such as adhesion to filters followed by washing, adhesion to plastic followed by washing, or centrifugation of the cell membranes. Viable cells could also be used to screen for the effects of drugs on cytokine mediated functions, e.g., STAT4 signaling and others. Some detection methods allow for elimination of a separation step, e.g., a proximity sensitive detection system.

The broad scope of this invention is best understood with reference to the following examples, which are not intended to limit the inventions to the specific embodiments.

## EXAMPLES

### I. General Methods

Some of the standard methods are described or referenced, e.g., in Maniatis, et al. (1982) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) *Molecular Cloning: A Laboratory Manual*, (2d ed.), vols. 1-3, CSH Press, NY; or Ausubel, et al. (1987 and Supplements) *Current Protocols in Molecular Biology*, Greene/Wiley, New York. Methods for protein purification include such methods as ammonium sulfate precipitation, column chromatography, electrophoresis, centrifugation, crystallization, and others. See, e.g., Ausubel, et al. (1987 and periodic supplements); Coligan, et al. (ed. 1996) and periodic supplements, *Current Protocols In Protein Science* Greene/Wiley, New York; Deutscher (1990) "Guide to Protein Purification" in *Methods in Enzymology*, vol. 182, and other volumes in this series; and manufacturer's

literature on use of protein purification products, e.g., Pharmacia, Piscataway, N.J., or Bio-Rad, Richmond, CA. Combination with recombinant techniques allow fusion to appropriate segments, e.g., to a FLAG sequence or an equivalent which can be fused via a protease-removable sequence. See, e.g., Hochuli (1990) "Purification of Recombinant Proteins with Metal Chelate Absorbent" in Setlow (ed.) Genetic Engineering, Principle and Methods 12:87-98, Plenum Press, N.Y.; and Crowe, et al. (1992) QIAexpress: The High Level Expression & Protein Purification System QUIAGEN, Inc., Chatsworth, CA.

Computer sequence analysis is performed, e.g., using available software programs, including those from the GCG (U. Wisconsin) and GenBank sources. Public sequence databases were also used, e.g., from GenBank and others.

Many techniques applicable to IL-10 receptors may be applied to the DCRS5, as described, e.g., in U.S. Patent No. 5,789,192 (IL-10 receptor), which is incorporated herein by reference.

## II. Functional cloning

It was observed that anti-hIL-12R $\beta$ 1 antibody blocked responses of human T cells to p40/IL-B30, and the p40/IL-B30 bound to IL-12R $\beta$ 1. This suggested that IL-12R $\beta$ 1 was one subunit of the receptor complex for p40/IL-B30.

A mouse T cell population was identified which responded to p40/IL-B30 but not to IL-12, and another population which responded to IL-12 but not p40/IL-B30. In addition, it was observed that Ba/F3 cells expressing recombinant mIL-12R $\beta$ 1 and mIL-12R $\beta$ 2 responded to IL-12, but not to p40/IL-B30. These results collectively indicated that the receptor complex for p40/IL-B30 contained the IL-12R $\beta$ 1 and at least one other subunit which was not IL-12R $\beta$ 2. Accordingly an expression cloning strategy was devised to isolate this second receptor component.

A cDNA library was prepared from mRNA isolated from Kit225 cells, an IL-2-dependent human T cell line which responds to both IL-12 and p40/IL-B30. The cDNA library was made using a retroviral expression vector, pMX. Ba/F3 cells expressing recombinant hIL-12R $\beta$ 1 were infected with this cDNA library, allowed to recover for 3-4 days in IL-3, then washed and plated at ~15,000 cells/well in 96 well plates in medium containing 50 ng/ml hyper-hp40/hIL-B30. See WO 01/18051. Cultures were supplemented every ~5 days with additional hyper-hp40/hIL-B30. After approximately two weeks 5-10% of the wells exhibited cell growth. Cells were recovered from each well, expanded individually in larger cultures in hyper-hp40/hIL-B30, and tested for growth dependence on hyper-hp40/hIL-B30.

Cells which were p40/IL-B30-dependent for growth were analyzed by PCR for retroviral cDNA inserts. Out of more than 40 isolates analyzed, all but one contained cDNAs

encoding the novel receptor DCRS5. This candidate human cDNA was cloned in an expression vector and transfected into Ba/F3 cells expressing hIL-12R $\beta$ 1. These cells became responsive to p40/IL-B30; thus we concluded that the novel cDNA encoded the desired DCRS5, functionally an IL-B30 receptor subunit.

5

### III. Features of full-length DCRS5; Chromosomal localization

The cytoplasmic domain of DCRS5 is not overall closely related to other cytokine receptor cytoplasmic domains, a common observation in this family of molecules. The cytoplasmic domain contains seven tyr residues, at least three of which are part of recognizable SH2-binding motifs: YEDI, YKPQ, and YFPQ. The YEDI motif is similar to identified binding sites for the tyrosine phosphatase shp2. The latter two motifs are very similar to sequences known to bind Stat1/Stat3, or Stat3, respectively. The YKPQ motif, together with nearby flanking sequences, also resembles to a degree the motifs in Stat4 and IL-12R $\beta$ 2 which are known to bind Stat1-3. This is consistent with preliminary data suggesting that p40/IL-B30, like IL-12, activates Stat4.

10  
15

PCR primers derived from the DCRS5 sequence are used to probe a human cDNA library. Sequences may be derived, e.g., from Table 1, preferably those adjacent the ends of sequences. Full length cDNAs for primate, rodent, or other species DCRS5 are cloned, e.g., by DNA hybridization screening of  $\lambda$ gt10 phage. PCR reactions are conducted using *T. aquaticus* Taqplus DNA polymerase (Stratagene) under appropriate conditions.

20

Chromosome spreads are prepared. In situ hybridization is performed on chromosome preparations obtained from phytohemagglutinin-stimulated human lymphocytes cultured for 72 h. 5-bromodeoxyuridine was added for the final seven hours of culture (60  $\mu$ g/ml of medium), to ensure a posthybridization chromosomal banding of good quality.

25

A PCR fragment, amplified with the help of primers, is cloned into an appropriate vector. The vector is labeled by nick-translation with  $^3$ H. The radiolabeled probe is hybridized to metaphase spreads at final concentration of 200 ng/ml of hybridization solution as described in Mattei, et al. (1985) Hum. Genet. 69:327-331.

30

After coating with nuclear track emulsion (KODAK NTB2), slides are exposed. To avoid any slipping of silver grains during the banding procedure, chromosome spreads are first stained with buffered Giemsa solution and metaphase photographed. R-banding is then performed by the fluorochrome-photolysis-Giemsa (FPG) method and metaphases rephotographed before analysis.

Similar appropriate methods are used for other species.

#### IV. Localization of DCRS5 mRNA

Human multiple tissue (Cat# 1, 2) and cancer cell line blots (Cat# 7757-1), containing approximately 2  $\mu$ g of poly (A) <sup>+</sup> RNA per lane, are purchased from Clontech (Palo Alto, CA).

5 Probes are radiolabeled with [ $\alpha$ -<sup>32</sup>P] dATP, e.g., using the Amersham Rediprime random primer labelling kit (RPN1633). Prehybridization and hybridizations are performed, e.g., at 65° C in 0.5 M Na<sub>2</sub>HPO<sub>4</sub>, 7% SDS, 0.5 M EDTA (pH 8.0). High stringency washes are conducted, e.g., at 65° C with two initial washes in 2 x SSC, 0.1% SDS for 40 min followed by a subsequent wash in 0.1 x SSC, 0.1% SDS for 20 min. Membranes are then exposed at -70° C  
10 to X-Ray film (Kodak) in the presence of intensifying screens. More detailed studies by cDNA library Southernblots are performed with selected appropriate human DCRS5 clones to examine their expression in hemopoietic or other cell subsets.

Alternatively, two appropriate primers are selected from Table 1. RT-PCR is used on an appropriate mRNA sample selected for the presence of message to produce a cDNA, e.g., a  
15 sample which expresses the gene.

Full length clones may be isolated by hybridization of cDNA libraries from appropriate tissues pre-selected by PCR signal. Northern blots can be performed.

Message for genes encoding DCRS5 will be assayed by appropriate technology, e.g., PCR, immunoassay, hybridization, or otherwise. Tissue and organ cDNA preparations are  
20 available, e.g., from Clontech, Mountain View, CA. Identification of sources of natural expression are useful, as described. And the identification of the functional receptor subunit pairing allows for prediction of what cells express the combination of receptor subunits which will result in a physiological responsiveness to each of the cytokine ligands.

For mouse distribution, e.g., Southern Analysis can be performed: DNA (5  $\mu$ g) from a  
25 primary amplified cDNA library was digested with appropriate restriction enzymes to release the inserts, run on a 1% agarose gel and transferred to a nylon membrane (Schleicher and Schuell, Keene, NY).

Samples for mouse mRNA isolation may include: resting mouse fibroblastic L cell line (C200); Braf:ER (Braf fusion to estrogen receptor) transfected cells, control (C201); T cells,  
30 TH1 polarized (Mell4 bright, CD4<sup>+</sup> cells from spleen, polarized for 7 days with IFN- $\gamma$  and anti IL-4; T200); T cells, TH2 polarized (Mell4 bright, CD4<sup>+</sup> cells from spleen, polarized for 7 days with IL-4 and anti-IFN- $\gamma$ ; T201); T cells, highly TH1 polarized (see Openshaw, et al. (1995) J. Exp. Med. 182:1357-1367; activated with anti-CD3 for 2, 6, 16 h pooled; T202); T cells, highly TH2 polarized (see Openshaw, et al. (1995) J. Exp. Med. 182:1357-1367;  
35 activated with anti-CD3 for 2, 6, 16 h pooled; T203); CD44-CD25<sup>+</sup> pre T cells, sorted from thymus (T204); TH1 T cell clone D1.1, resting for 3 weeks after last stimulation with antigen

- (T205); TH1 T cell clone D1.1, 10 µg/ml ConA stimulated 15 h (T206); TH2 T cell clone CDC35, resting for 3 weeks after last stimulation with antigen (T207); TH2 T cell clone CDC35, 10 µg/ml ConA stimulated 15 h (T208); Mell4+ naïve T cells from spleen, resting (T209); Mell4+ T cells, polarized to Th1 with IFN-γ/IL-12/anti-IL-4 for 6, 12, 24 h pooled (T210); Mell4+ T cells, polarized to Th2 with IL-4/anti-IFN-γ for 6, 13, 24 h pooled (T211); unstimulated mature B cell leukemia cell line A20 (B200); unstimulated B cell line CH12 (B201); unstimulated large B cells from spleen (B202); B cells from total spleen, LPS activated (B203); metrizamide enriched dendritic cells from spleen, resting (D200); dendritic cells from bone marrow, resting (D201); monocyte cell line RAW 264.7 activated with LPS 4 h (M200); bone-marrow macrophages derived with GM and M-CSF (M201); macrophage cell line J774, resting (M202); macrophage cell line J774 + LPS + anti-IL-10 at 0.5, 1, 3, 6, 12 h pooled (M203); macrophage cell line J774 + LPS + IL-10 at 0.5, 1, 3, 5, 12 h pooled (M204); aerosol challenged mouse lung tissue, Th2 primers, aerosol OVA challenge 7, 14, 23 h pooled (see Garlisi, et al. (1995) *Clinical Immunology and Immunopathology* 75:75-83; X206);
- 15 Nippostrongylus-infected lung tissue (see Coffman, et al. (1989) *Science* 245:308-310; X200); total adult lung, normal (O200); total lung, rag-1 (see Schwarz, et al. (1993) *Immunodeficiency* 4:249-252; O205); IL-10 K.O. spleen (see Kuhn, et al. (1991) *Cell* 75:263-274; X201); total adult spleen, normal (O201); total spleen, rag-1 (O207); IL-10 K.O. Peyer's patches (O202); total Peyer's patches, normal (O210); IL-10 K.O. mesenteric lymph nodes (X203); total
- 20 mesenteric lymph nodes, normal (O211); IL-10 K.O. colon (X203); total colon, normal (O212); NOD mouse pancreas (see Makino, et al. (1980) *Jikken Dobutsu* 29:1-13; X205); total thymus, rag-1 (O208); total kidney, rag-1 (O209); total heart, rag-1 (O202); total brain, rag-1 (O203); total testes, rag-1 (O204); total liver, rag-1 (O206); rat normal joint tissue (O300); and rat arthritic joint tissue (X300).
- 25 Samples for human mRNA isolation may include: peripheral blood mononuclear cells (monocytes, T cells, NK cells, granulocytes, B cells), resting (T100); peripheral blood mononuclear cells, activated with anti-CD3 for 2, 6, 12 h pooled (T101); T cell, TH0 clone Mot 72, resting (T102); T cell, TH0 clone Mot 72, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T103); T cell, TH0 clone Mot 72, anergic treated with specific peptide for 2,
- 30 7, 12 h pooled (T104); T cell, TH1 clone HY06, resting (T107); T cell, TH1 clone HY06, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T108); T cell, TH1 clone HY06, anergic treated with specific peptide for 2, 6, 12 h pooled (T109); T cell, TH2 clone HY935, resting (T110); T cell, TH2 clone HY935, activated with anti-CD28 and anti-CD3 for 2, 7, 12 h pooled (T111); T cells CD4+CD45RO- T cells polarized 27 days in anti-CD28, IL-4, and anti
- 35 IFN-γ, TH2 polarized, activated with anti-CD3 and anti-CD28 4 h (T116); T cell tumor lines Jurkat and Hut78, resting (T117); T cell clones, pooled AD130.2, Tc783.12, Tc783.13,



Tc783.58, Tc782.69, resting (T118); T cell random  $\gamma\delta$  T cell clones, resting (T119); Splenocytes, resting (B100); Splenocytes, activated with anti-CD40 and IL-4 (B101); B cell EBV lines pooled WT49, RSB, JY, CVIR, 721.221, RM3, HSY, resting (B102); B cell line JY, activated with PMA and ionomycin for 1, 6 h pooled (B103); NK 20 clones pooled, resting (K100); NK 20 clones pooled, activated with PMA and ionomycin for 6 h (K101); NKL clone, derived from peripheral blood of LGL leukemia patient, IL-2 treated (K106); NK cytotoxic clone 640-A30-1, resting (K107); hematopoietic precursor line TF1, activated with PMA and ionomycin for 1, 6 h pooled (C100); U937 premonocytic line, resting (M100); U937 premonocytic line, activated with PMA and ionomycin for 1, 6 h pooled (M101); elutriated monocytes, activated with LPS, IFN $\gamma$ , anti-IL-10 for 1, 2, 6, 12, 24 h pooled (M102); elutriated monocytes, activated with LPS, IFN $\gamma$ , IL-10 for 1, 2, 6, 12, 24 h pooled (M103); elutriated monocytes, activated with LPS, IFN $\gamma$ , anti-IL-10 for 4, 16 h pooled (M106); elutriated monocytes, activated with LPS, IFN $\gamma$ , IL-10 for 4, 16 h pooled (M107); elutriated monocytes, activated LPS for 1 h (M108); elutriated monocytes, activated LPS for 6 h (M109); DC 70% CD1a+, from CD34+ GM-CSF, TNF $\alpha$  12 days, resting (D101); DC 70% CD1a+, from CD34+ GM-CSF, TNF $\alpha$  12 days, activated with PMA and ionomycin for 1 hr (D102); DC 70% CD1a+, from CD34+ GM-CSF, TNF $\alpha$  12 days, activated with PMA and ionomycin for 6 hr (D103); DC 95% CD1a+, from CD34+ GM-CSF, TNF $\alpha$  12 days FACS sorted, activated with PMA and ionomycin for 1, 6 hr pooled (D104); DC 95% CD14+, ex CD34+ GM-CSF, TNF $\alpha$  12 days FACS sorted, activated with PMA and ionomycin 1, 6 hr pooled (D105); DC CD1a+ CD86+, from CD34+ GM-CSF, TNF $\alpha$  12 days FAC sorted, activated with PMA and ionomycin for 1, 6 h pooled (K106); DC from monocytes GM-CSF, IL-4 5 days, resting (D107); DC from monocytes GM-CSF, IL-4 5 days, resting (D108); DC from monocytes GM-CSF, IL-4 5 days, activated LPS 4, 16 h pooled (D109); DC from monocytes GM-CSF, IL-4 5 days, activated TNF $\alpha$ , monocyte supe for 4, 16 h pooled (D110); leiomyoma L11 benign tumor (X101); normal myometrium M5 (O115); malignant leiomyosarcoma GS1 (X103); lung fibroblast sarcoma line MRC5, activated with PMA and ionomycin for 1, 6 h pooled (C101); kidney epithelial carcinoma cell line CHA, activated with PMA and ionomycin for 1, 6 h pooled (C102); kidney fetal 28 wk male (O100); lung fetal 28 wk male (O101); liver fetal 28 wk male (O102); heart fetal 28 wk male (O103); brain fetal 28 wk male (O104); gallbladder fetal 28 wk male (O106); small intestine fetal 28 wk male (O107); adipose tissue fetal 28 wk male (O108); ovary fetal 25 wk female (O109); uterus fetal 25 wk female (O110); testes fetal 28 wk male (O111); spleen fetal 28 wk male (O112); adult placenta 28 wk (O113); and tonsil inflamed, from 12 year old (X100).

Similar samples may isolated in other species for evaluation.

## V. Cloning of species counterparts of DCRS5

Various strategies are used to obtain species counterparts of the DCRS5, preferably from other primates or rodents. One method is by cross hybridization using closely related species DNA probes. It may be useful to go into evolutionarily similar species as intermediate steps. Another method is by using specific PCR primers based on the identification of blocks of similarity or difference between genes, e.g., areas of highly conserved or nonconserved polypeptide or nucleotide sequence.

Database searches may identify similar sequences and allow production of appropriate probes.

## VI. Production of mammalian DCRS5 protein

An appropriate, e.g., GST, fusion construct is engineered for expression, e.g., in *E. coli*. For example, a mouse IGIF pGex plasmid is constructed and transformed into *E. coli*. Freshly transformed cells are grown, e.g., in LB medium containing 50 µg/ml ampicillin and induced with IPTG (Sigma, St. Louis, MO). After overnight induction, the bacteria are harvested and the pellets containing the DCRS5 protein are isolated. The pellets are homogenized, e.g., in TE buffer (50 mM Tris-base pH 8.0, 10 mM EDTA and 2 mM pefabloc) in 2 liters. This material is passed through a microfluidizer (Microfluidics, Newton, MA) three times. The fluidized supernatant is spun down on a Sorvall GS-3 rotor for 1 h at 13,000 rpm. The resulting supernatant containing the cytokine receptor protein is filtered and passed over a glutathione-SEPHAROSE column equilibrated in 50 mM Tris-base pH 8.0. The fractions containing the DCRS5-GST fusion protein are pooled and cleaved, e.g., with thrombin (Enzyme Research Laboratories, Inc., South Bend, IN). The cleaved pool is then passed over a Q-SEPHAROSE column equilibrated in 50 mM Tris-base. Fractions containing DCRS5 are pooled and diluted in cold distilled H<sub>2</sub>O, to lower the conductivity, and passed back over a fresh Q-Sepharose column, alone or in succession with an immunoaffinity antibody column. Fractions containing the DCRS5 protein are pooled, aliquoted, and stored in the -70° C freezer.

Comparison of the CD spectrum with cytokine receptor protein may suggest that the protein is correctly folded. See Hazuda, et al. (1969) *J. Biol. Chem.* 264:1689-1693.

## VII. Preparation of antibodies specific for DCRS5

Inbred Balb/c mice are immunized intraperitoneally with recombinant forms of the protein, e.g., purified DCRS5 or stable transfected NIH-3T3 cells. Animals are boosted at appropriate time points with protein, with or without additional adjuvant, to further stimulate antibody production. Serum is collected, or hybridomas produced with harvested spleens.

Alternatively, Balb/c mice are immunized with cells transformed with the gene or fragments thereof, either endogenous or exogenous cells, or with isolated membranes enriched for expression of the antigen. Serum is collected at the appropriate time, typically after numerous further administrations. Various gene therapy techniques may be useful, e.g., in  
5 producing protein in situ, for generating an immune response. Serum or antibody preparations may be cross-absorbed or immunoselected to prepare substantially purified antibodies of defined specificity and high affinity.

Monoclonal antibodies may be made. For example, splenocytes are fused with an appropriate fusion partner and hybridomas are selected in growth medium by standard  
10 procedures. Hybridoma supernatants are screened for the presence of antibodies which bind to the DCRS5, e.g., by ELISA or other assay. Antibodies which specifically recognize specific DCRS5 embodiments may also be selected or prepared.

In another method, synthetic peptides or purified protein are presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (ed. 1991) Current  
15 Protocols in Immunology Wiley/Greene; and Harlow and Lane (1989) Antibodies: A Laboratory Manual Cold Spring Harbor Press. In appropriate situations, the binding reagent is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to a substrate for panning methods. Nucleic acids may also be introduced into cells in an animal to produce the antigen, which serves to elicit an immune response. See, e.g., Wang, et al. (1993) Proc.  
20 Nat'l. Acad. Sci. 90:4156-4160; Barry, et al. (1994) BioTechniques 16:616-619; and Xiang, et al. (1995) Immunity 2: 129-135.

#### VIII. Production of fusion proteins with DCRS5

Various fusion constructs are made with DCRS5, including embodiments combining  
25 such with IL-12R $\beta$ 1 sequence. A portion of the appropriate gene is fused to an epitope tag, e.g., a FLAG tag, or to a two hybrid system construct. See, e.g., Fields and Song (1989) Nature 340:245-246.

The epitope tag may be used in an expression cloning procedure with detection with anti-FLAG antibodies to detect a binding partner, e.g., ligand for the respective cytokine  
30 receptor. The two hybrid system may also be used to isolate proteins which specifically bind to DCRS5.

#### IX. Structure activity relationship

Information on the criticality of particular residues is determined using standard  
35 procedures and analysis. Standard mutagenesis analysis is performed, e.g., by generating many different variants at determined positions, e.g., at the positions identified above, and evaluating

biological activities of the variants. This may be performed to the extent of determining positions which modify activity, or to focus on specific positions to determine the residues which can be substituted to either retain, block, or modulate biological activity.

Alternatively, analysis of natural variants can indicate what positions tolerate natural  
5 mutations. This may result from populational analysis of variation among individuals, or across strains or species. Samples from selected individuals are analyzed, e.g., by PCR analysis and sequencing. This allows evaluation of population polymorphisms.

#### X. Coexpression of DCRS5 and IL-12R $\beta$ 1

10 A vector, or vectors, encoding the respective genes may be transfected into a cell. Preferably, such vector will have selection markers to identify which cells have successfully been transformed. Coexpression of the two genes will allow the gene products to properly associate to form active receptor complexes.

Alternatively, use of methods causing association of functional dimers are available.  
15 See, e.g., O'Shea, et al. (1989) Science 245:646-648; Kostelny, et al. (1992) J. Immunol. 148:1547-1553; and Patel, et al. (1996) J. Biol. Chem. 271:30386-30391. Expression of extracellular domains, and physical association, e.g., driven by Fos/Jun leucine zipper affinity, will result in ligand binding constructs which should act as binding compounds for diagnostic or therapeutic uses.

20 All citations herein are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific  
25 embodiments described herein are offered by way of example only, and the invention is to be limited by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled; and the invention is not to be limited by the specific embodiments that have been presented herein by way of example.